

# Anti-ACLY Antibody

Rabbit polyclonal antibody to ACLY  
Catalog # AP60530

## Product Information

Application	WB, E, IF/IC, IHC
Primary Accession	<a href="#">P53396</a>
Other Accession	<a href="#">Q91V92</a>
Reactivity	Human, Mouse, Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	120839

## Additional Information


Gene ID	47
Other Names	ATP-citrate synthase; ATP-citrate (pro-S-)-lyase; ACL; Citrate cleavage enzyme
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ACLY. The exact sequence is proprietary.
Dilution	WB~~E (1/5000 - 1/10000), WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500) E~~N/A IF/IC~~N/A IHC~~E (1/5000 - 1/10000), WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information


Name	ACLY
Function	Catalyzes the cleavage of citrate into oxaloacetate and acetyl-CoA, the latter serving as common substrate in multiple biochemical reactions in protein, carbohydrate and lipid metabolism.
Cellular Location	Cytoplasm, cytosol.

## Background


KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ACLY. The exact sequence is proprietary.



Western blot analysis of ACLY expression in C6 (A) whole cell lysates.



Immunohistochemical analysis of ACLY staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ACLY staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.