

Anti-Kv7.5 Antibody

Rabbit polyclonal antibody to Kv7.5 Catalog # AP60906

Product Information

ApplicationWBPrimary AccessionQ9NR82Other AccessionQ9JK45

Reactivity Human, Mouse, Monkey

Host Rabbit
Clonality Polyclonal
Calculated MW 102179

Additional Information

Gene ID 56479

Other Names Potassium voltage-gated channel subfamily KQT member 5; KQT-like 5;

Potassium channel subunit alpha KvLQT5; Voltage-gated potassium channel

subunit Kv7.5

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human Kv7.5. The exact sequence is proprietary.

Dilution WB~~WB (1/500 - 1/1000)

Format Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.09% (W/V) sodium azide.

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name KCNQ5 (<u>HGNC:6299</u>)

Function Pore-forming subunit of the voltage-gated potassium (Kv) channel broadly

expressed in brain and involved in the regulation of neuronal excitability

(PubMed: 10787416, PubMed: 10816588, PubMed: 11159685,

PubMed:<u>28669405</u>). Associates with KCNQ3/Kv7.3 pore- forming subunit to form a potassium channel which contributes to M-type current, a slowly activating and deactivating potassium conductance which plays a critical role

in determining the subthreshold electrical excitability of neurons

(PubMed:10816588, PubMed:11159685). Contributes, with other potassium channels, to the molecular diversity of a heterogeneous population of M-channels, varying in kinetic and pharmacological properties, which underlie this physiologically important current (PubMed:10816588). Also forms a functional channel with KCNQ1/Kv7.1 subunit that may contribute to

vasoconstriction and hypertension (PubMed:24855057). Channel may be selectively permeable in vitro to other cations besides potassium, in decreasing order of affinity K(+) = Rb(+) > Cs(+) > Na(+) (PubMed:10816588). Similar to the native M-channel, KCNQ3-KCNQ5 potassium channel is suppressed by activation of the muscarinic acetylcholine receptor CHRM1 (PubMed:10816588).

Cellular Location

Cell membrane; Multi-pass membrane protein


Tissue Location

Strongly expressed in brain and skeletal muscle (PubMed:10787416, PubMed:10816588). In brain, expressed in cerebral cortex, occipital pole, frontal lobe and temporal lobe. Lower levels in hippocampus and putamen. Low to undetectable levels in medulla, cerebellum and thalamus (PubMed:10787416, PubMed:10816588)

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Kv7.5. The exact sequence is proprietary.

Images



Western blot analysis of Kv7.5 expression in HEK293T (A), COS7 (B), C6 (C), CT26 (D), MG63 (E) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.