

Anti-PINK1 (pS402) Antibody

Rabbit polyclonal antibody to PINK1 (pS402)
Catalog # AP61087

Product Information

Application	WB
Primary Accession	Q9BXM7
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62769

Additional Information

Gene ID	65018
Other Names	Serine/threonine-protein kinase PINK1 mitochondrial; BRPK; PTEN-induced putative kinase protein 1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PINK1. The exact sequence is proprietary.
Dilution	WB--WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information

Name	PINK1
Function	Serine/threonine-protein kinase which acts as a sensor of mitochondrial damage and protects against mitochondrial dysfunction during cellular stress (PubMed: 40080546). It phosphorylates mitochondrial proteins to coordinate mitochondrial quality control mechanisms that remove and replace dysfunctional mitochondrial components (PubMed: 14607334 , PubMed: 15087508 , PubMed: 18443288 , PubMed: 18957282 , PubMed: 19229105 , PubMed: 19966284 , PubMed: 20404107 , PubMed: 20547144 , PubMed: 20798600 , PubMed: 22396657 , PubMed: 23620051 , PubMed: 23754282 , PubMed: 23933751 , PubMed: 24660806 , PubMed: 24751536 , PubMed: 24784582 , PubMed: 24896179 , PubMed: 24898855 , PubMed: 25527291 , PubMed: 32484300). In healthy mitochondria, PINK1 is translocated across the mitochondrial outer membrane (MOM) via the translocase of the outer membrane (TOM) complex, and inserted into the mitochondrial inner

membrane (MIM) via the translocase of the inner membrane (TIM23) complex where it is cleaved and released into the cytosol (PubMed:[40080546](#)). Depending on the severity of mitochondrial damage, activity ranges from preventing apoptosis and stimulating mitochondrial biogenesis to eliminating severely damaged mitochondria via PINK1-PRKN- dependent mitophagy (PubMed:[14607334](#), PubMed:[15087508](#), PubMed:[18443288](#), PubMed:[19966284](#), PubMed:[20404107](#), PubMed:[20798600](#), PubMed:[22396657](#), PubMed:[23620051](#), PubMed:[23933751](#), PubMed:[24898855](#), PubMed:[32047033](#), PubMed:[32484300](#)). When cellular stress results in irreversible mitochondrial damage, PINK1 accumulates at the outer mitochondrial membrane (OMM) where it phosphorylates pre-existing polyubiquitin chains at 'Ser-65', recruits PRKN from the cytosol to the OMM and activates PRKN by phosphorylation at 'Ser-65'; activated PRKN then ubiquitinates VDAC1 and other OMM proteins to initiate mitophagy (PubMed:[14607334](#), PubMed:[15087508](#), PubMed:[19966284](#), PubMed:[20404107](#), PubMed:[20798600](#), PubMed:[23754282](#), PubMed:[23933751](#), PubMed:[24660806](#), PubMed:[24751536](#), PubMed:[24784582](#), PubMed:[25474007](#), PubMed:[25527291](#), PubMed:[32047033](#), PubMed:[40080546](#)). The PINK1-PRKN pathway also promotes fission of damaged mitochondria through phosphorylation and PRKN- dependent degradation of mitochondrial proteins involved in fission such as MFN2 (PubMed:[18443288](#), PubMed:[23620051](#), PubMed:[24898855](#)). This prevents the refusion of unhealthy mitochondria with the mitochondrial network or initiates mitochondrial fragmentation facilitating their later engulfment by autophagosomes (PubMed:[18443288](#), PubMed:[23620051](#)). Also promotes mitochondrial fission independently of PRKN and ATG7-mediated mitophagy, via the phosphorylation and activation of DNM1L (PubMed:[18443288](#), PubMed:[32484300](#)). Regulates motility of damaged mitochondria by promoting the ubiquitination and subsequent degradation of MIRO1 and MIRO2; in motor neurons, this likely inhibits mitochondrial intracellular anterograde transport along the axons which probably increases the chance of the mitochondria undergoing mitophagy in the soma (PubMed:[22396657](#)). Required for ubiquinone reduction by mitochondrial complex I by mediating phosphorylation of complex I subunit NDUFA10 (By similarity). Phosphorylates LETM1, positively regulating its mitochondrial calcium transport activity (PubMed:[29123128](#)).

Cellular Location

Mitochondrion outer membrane Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q99MQ3}. Cytoplasm, cytosol Note=Localizes mostly in mitochondrion and the two smaller proteolytic processed fragments localize mainly in cytosol (PubMed:[19229105](#)). In healthy mitochondria, PINK1 is translocated across the mitochondrial membranes (PubMed:[40080546](#)). Upon mitochondrial membrane depolarization following damage, PINK1 import is stalled, which induces its accumulation in the outer mitochondrial membrane and the activation of its kinase activity (PubMed:[18957282](#), PubMed:[40080546](#))


Tissue Location

Highly expressed in heart, skeletal muscle and testis, and at lower levels in brain, placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Present in the embryonic testis from an early stage of development

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PINK1. The exact sequence is proprietary.

Images



Western blot analysis of PINK1 (pS402) expression in MCF7 (A), HCT116 (B) whole cell lysates.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.