

Anti-Caspase 7 p20 Antibody

Rabbit polyclonal antibody to Caspase 7 p20

Catalog # AP61403

Product Information

Application	WB, IHC
Primary Accession	P55210
Other Accession	P97864
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34277

Additional Information

Gene ID	840
Other Names	MCH3; Caspase-7; CASP-7; Apoptotic protease Mch-3; CMH-1; ICE-like apoptotic protease 3; ICE-LAP3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Caspase 7 p20. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000), IHC (1/50 - 1/200) IHC~~WB (1/500 - 1/1000), IHC (1/50 - 1/200)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CASP7 {ECO:0000303 PubMed:9070923, ECO:0000312 HGNC:HGNC:1508}
Function	Thiol protease involved in different programmed cell death processes, such as apoptosis, pyroptosis or granzyme-mediated programmed cell death, by proteolytically cleaving target proteins (PubMed: 11257230 , PubMed: 11257231 , PubMed: 11701129 , PubMed: 15314233 , PubMed: 16916640 , PubMed: 17646170 , PubMed: 18723680 , PubMed: 19581639 , PubMed: 8521391 , PubMed: 8567622 , PubMed: 8576161 , PubMed: 9070923). Has a marked preference for Asp-Glu-Val-Asp (DEVD) consensus sequences, with some plasticity for alternate non-canonical sequences (PubMed: 12824163 , PubMed: 15314233 , PubMed: 17697120 , PubMed: 19581639 , PubMed: 20566630 , PubMed: 23650375 , PubMed: 23897474 , PubMed: 27032039). Its involvement in the different programmed cell death processes is probably determined by upstream

proteases that activate CASP7 (By similarity). Acts as an effector caspase involved in the execution phase of apoptosis: following cleavage and activation by initiator caspases (CASP8, CASP9 and/or CASP10), mediates execution of apoptosis by catalyzing cleavage of proteins, such as CLSPN, PARP1, PTGES3 and YY1 (PubMed:[10497198](#), PubMed:[16123041](#), PubMed:[16374543](#), PubMed:[16916640](#), PubMed:[18723680](#), PubMed:[20566630](#), PubMed:[21555521](#), PubMed:[22184066](#), PubMed:[22451931](#), PubMed:[27889207](#), PubMed:[28863261](#), PubMed:[31586028](#), PubMed:[34156061](#), PubMed:[35338844](#), PubMed:[35446120](#)). Compared to CASP3, acts as a minor executioner caspase and cleaves a limited set of target proteins (PubMed:[18723680](#)). Acts as a key regulator of the inflammatory response in response to bacterial infection by catalyzing cleavage and activation of the sphingomyelin phosphodiesterase SMPD1 in the extracellular milieu, thereby promoting membrane repair (PubMed:[21157428](#)). Regulates pyroptosis in intestinal epithelial cells: cleaved and activated by CASP1 in response to S.typhimurium infection, promoting its secretion to the extracellular milieu, where it catalyzes activation of SMPD1, generating ceramides that repair membranes and counteract the action of gasdermin-D (GSDMD) pores (By similarity). Regulates granzyme-mediated programmed cell death in hepatocytes: cleaved and activated by granzyme B (GZMB) in response to bacterial infection, promoting its secretion to the extracellular milieu, where it catalyzes activation of SMPD1, generating ceramides that repair membranes and counteract the action of perforin (PRF1) pores (By similarity). Following cleavage by CASP1 in response to inflammasome activation, catalyzes processing and inactivation of PARP1, alleviating the transcription repressor activity of PARP1 (PubMed:[22464733](#)). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (By similarity). Cleaves and activates sterol regulatory element binding proteins (SREBPs) (PubMed:[8643593](#)). Cleaves phospholipid scramblase proteins XKR4, XKR8 and XKR9 (By similarity). In case of infection, catalyzes cleavage of Kaposi sarcoma-associated herpesvirus protein ORF57, thereby preventing expression of viral lytic genes (PubMed:[20159985](#)). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:[36758104](#), PubMed:[36758106](#)).

Cellular Location

Cytoplasm, cytosol. Nucleus. Secreted, extracellular space {ECO:0000250|UniProtKB:P97864}. Note=Following cleavage and activation by CASP1 or granzyme B (GZMB), secreted into the extracellular milieu by passing through the gasdermin-D (GSDMD) pores or perforin (PRF1) pore, respectively {ECO:0000250|UniProtKB:P97864}

Tissue Location



Highly expressed in lung, skeletal muscle, liver, kidney, spleen and heart, and moderately in testis. No expression in the brain.

Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Caspase 7 p20. The exact sequence is proprietary.

Images

Western blot analysis of Caspase 7 p20 expression in rat lung (A), rat liver (B) whole cell lysates.



Immunohistochemical analysis of Caspase 7 p20 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.