

Presenilin 1 (PSEN1) Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6231A

Product Information

Application	WB, IHC-P, FC, IF, E
Primary Accession	P49768
Other Accession	P97887 , P49769 , Q8HXW5
Reactivity	Human, Rat, Mouse
Predicted	Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52668
Antigen Region	330-359

Additional Information

Gene ID	5663
Other Names	Presenilin-1, PS-1, 3423-, Protein S182, Presenilin-1 NTF subunit, Presenilin-1 CTF subunit, Presenilin-1 CTF12, PS1-CTF12, PSEN1, AD3, PS1, PSNL1
Target/Specificity	This Presenilin 1 (PSEN1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 330-359 amino acids from the C-terminal region of human Presenilin 1 (PSEN1).
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Presenilin 1 (PSEN1) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PSEN1 (HGNC:9508)
Synonyms	AD3, PS1, PSNL1


Function	Catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed: 10206644 , PubMed: 10545183 , PubMed: 10593990 , PubMed: 10811883 , PubMed: 10899933 , PubMed: 12679784 , PubMed: 12740439 , PubMed: 15274632 , PubMed: 20460383 , PubMed: 25043039 , PubMed: 26280335 , PubMed: 28269784 , PubMed: 30598546 , PubMed: 30630874). Requires the presence of the other members of the gamma-secretase complex for protease activity (PubMed: 15274632 , PubMed: 25043039 , PubMed: 26280335 , PubMed: 30598546 , PubMed: 30630874). Plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels (PubMed: 10593990 , PubMed: 10811883 , PubMed: 10899933 , PubMed: 9738936). Stimulates cell-cell adhesion via its interaction with CDH1; this stabilizes the complexes between CDH1 (E- cadherin) and its interaction partners CTNNB1 (beta-catenin), CTNND1 and JUP (gamma-catenin) (PubMed: 11953314). Under conditions of apoptosis or calcium influx, cleaves CDH1 (PubMed: 11953314). This promotes the disassembly of the complexes between CDH1 and CTNND1, JUP and CTNNB1, increases the pool of cytoplasmic CTNNB1, and thereby negatively regulates Wnt signaling (PubMed: 11953314 , PubMed: 9738936). Required for normal embryonic brain and skeleton development, and for normal angiogenesis (By similarity). Mediates the proteolytic cleavage of EphB2/CTF1 into EphB2/CTF2 (PubMed: 17428795 , PubMed: 28269784). The holoprotein functions as a calcium-leak channel that allows the passive movement of calcium from endoplasmic reticulum to cytosol and is therefore involved in calcium homeostasis (PubMed: 16959576 , PubMed: 25394380). Involved in the regulation of neurite outgrowth (PubMed: 15004326 , PubMed: 20460383). Is a regulator of presynaptic facilitation, spike transmission and synaptic vesicles replenishment in a process that depends on gamma-secretase activity. It acts through the control of SYT7 presynaptic expression (By similarity).
Cellular Location	Endoplasmic reticulum. Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic granule. Cell membrane; Multi-pass membrane protein. Cell projection, growth cone. Early endosome. Early endosome membrane; Multi-pass membrane protein. Cell projection, neuron projection. Cell projection, axon {ECO:0000250 UniProtKB:Q4JIM4}. Synapse {ECO:0000250 UniProtKB:Q4JIM4}. Note=Translocates with bound NOTCH1 from the endoplasmic reticulum and/or Golgi to the cell surface (PubMed:10593990). Colocalizes with CDH1/2 at sites of cell-cell contact. Colocalizes with CTNNB1 in the endoplasmic reticulum and the proximity of the plasma membrane (PubMed:9738936). Also present in azurophil granules of neutrophils (PubMed:11987239). Colocalizes with UBQLN1 in the cell membrane and in cytoplasmic juxtanuclear structures called aggresomes (PubMed:21143716). Also highly enriched in mitochondria-associated endoplasmic reticulum membrane contact site (By similarity). {ECO:0000250 UniProtKB:P49769, ECO:0000269 PubMed:10593990, ECO:0000269 PubMed:11987239, ECO:0000269 PubMed:21143716, ECO:0000269 PubMed:9738936}
Tissue Location	Detected in azurophile granules in neutrophils and in platelet cytoplasmic granules (at protein level) (PubMed:11987239) Expressed in a wide range of tissues including various regions of the brain, liver, spleen and lymph nodes (PubMed:7596406, PubMed:8574969, PubMed:8641442).
Background	

Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes.


References

Marambaud, P., et al., Cell 114(5):635-645 (2003). Kim, S.H., et al., J. Biol. Chem. 278(36):33992-34002 (2003). Miklossy, J., et al., Neurobiol. Aging 24(5):655-662 (2003). Cai, D., et al., J. Biol. Chem. 278(5):3446-3454 (2003). Godin, C., et al., Neuroreport 14(12):1613-1616 (2003).


Images



The Presenilin 1 (PSEN1) Antibody (C-term) (Cat.#AP6231a) is used in Western blot to detect PSEN1 in mouse kidney tissue lysate (lane 1) and HL60 cell lysate (lane 2) lysate.




Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Presenilin 1 (PSEN1) Antibody (C-term) (Cat. #AP6231a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Confocal immunofluorescent analysis of Presenilin 1 (PSEN1) Antibody (C-term)(Cat#AP6231a) with MDA-MB435 cell followed by Alexa Fluor 488-conjugated



goat anti-rabbit IgG (green).DAPI was used to stain the cell nuclear (blue).

Citations

- [Nicotine decreases beta-amyloid through regulating BACE1 transcription in SH-EP1- \$\alpha\$ 4 \$\beta\$ 2 nAChR-APP695 cells.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.