

Cleaved-Caspase-4 p20 (Q81) Polyclonal Antibody

Catalog # AP63103

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P49662
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43262

Additional Information

Gene ID	837
Other Names	CASP4; ICH2; Caspase-4; CASP-4; ICE(rel)-II; Protease ICH-2; Protease TX
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CASP4 {ECO:0000303 PubMed:15123740, ECO:0000312 HGNC:HGNC:1505}
Function	Inflammatory caspase that acts as the effector of the non- canonical inflammasome by mediating lipopolysaccharide (LPS)-induced pyroptosis (PubMed: 25119034 , PubMed: 26375003 , PubMed: 32109412 , PubMed: 34671164 , PubMed: 37001519 , PubMed: 37993712 , PubMed: 37993714). Also indirectly activates the NLRP3 and NLRP6 inflammasomes (PubMed: 23516580 , PubMed: 26375003 , PubMed: 32109412 , PubMed: 7797510). Acts as a thiol protease that cleaves a tetrapeptide after an Asp residue at position P1: catalyzes cleavage of CGAS, GSDMD and IL18 (PubMed: 15326478 , PubMed: 23516580 , PubMed: 26375003 , PubMed: 28314590 , PubMed: 32109412 , PubMed: 37993712 , PubMed: 37993714 , PubMed: 7797510). Effector of the non-canonical inflammasome independently of NLRP3 inflammasome and CASP1: the non-canonical inflammasome promotes pyroptosis through GSDMD cleavage without involving secretion of cytokine IL1B (PubMed: 25119034 , PubMed: 25121752 , PubMed: 26375003 , PubMed: 31268602 , PubMed: 32109412 , PubMed: 37993712 , PubMed: 37993714). In the non-canonical inflammasome, CASP4 is activated by direct binding to the lipid

A moiety of LPS without the need of an upstream sensor (PubMed:[25119034](#), PubMed:[25121752](#), PubMed:[29520027](#), PubMed:[32510692](#), PubMed:[32581219](#), PubMed:[37993712](#)). LPS-binding promotes CASP4 activation and CASP4-mediated cleavage of GSDMD and IL18, followed by IL18 secretion through the GSDMD pore, pyroptosis of infected cells and their extrusion into the gut lumen (PubMed:[25119034](#), PubMed:[25121752](#), PubMed:[37993712](#), PubMed:[37993714](#)). Also indirectly promotes secretion of mature cytokines (IL1A and HMGB1) downstream of GSDMD-mediated pyroptosis via activation of the NLRP3 and NLRP6 inflammasomes (PubMed:[26375003](#), PubMed:[32109412](#)). Involved in NLRP3-dependent CASP1 activation and IL1B secretion in response to non-canonical activators, such as UVB radiation or cholera enterotoxin (PubMed:[22246630](#), PubMed:[23516580](#), PubMed:[24879791](#), PubMed:[25964352](#), PubMed:[26173988](#), PubMed:[26174085](#), PubMed:[26508369](#)). Involved in NLRP6 inflammasome-dependent activation in response to lipoteichoic acid (LTA), a cell- wall component of Gram-positive bacteria, which leads to CASP1 activation and IL1B secretion (PubMed:[33377178](#)). Involved in LPS- induced IL6 secretion; this activity may not require caspase enzymatic activity (PubMed:[26508369](#)). The non-canonical inflammasome is required for innate immunity to cytosolic, but not vacuolar, bacteria (By similarity). Plays a crucial role in the restriction of *S.typhimurium* replication in colonic epithelial cells during infection (PubMed:[25121752](#), PubMed:[25964352](#)). Activation of the non-canonical inflammasome in brain endothelial cells can lead to excessive pyroptosis, leading to blood-brain barrier breakdown (By similarity). Pyroptosis limits bacterial replication, while cytokine secretion promotes the recruitment and activation of immune cells and triggers mucosal inflammation (PubMed:[25121752](#), PubMed:[25964352](#), PubMed:[26375003](#)). May also act as an activator of adaptive immunity in dendritic cells, following activation by oxidized phospholipid 1- palmitoyl-2-arachidonoyl-sn-glycero-3-phosphorylcholine, an oxidized phospholipid (oxPAPC) (By similarity). Involved in cell death induced by endoplasmic reticulum stress and by treatment with cytotoxic APP peptides found in Alzheimer's patient brains (PubMed:[15123740](#), PubMed:[22246630](#), PubMed:[23661706](#)). Cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP4 that recognizes and binds the Gasdermin-D, C- terminal (GSDMD-CT) part (PubMed:[32109412](#)). Catalyzes cleavage and maturation of IL18; IL18 processing also depends of the exosite interface on CASP4 (PubMed:[15326478](#), PubMed:[37993712](#), PubMed:[37993714](#)). In contrast, it does not directly process IL1B (PubMed:[7743998](#), PubMed:[7797510](#), PubMed:[7797592](#)). During non-canonical inflammasome activation, cuts CGAS and may play a role in the regulation of antiviral innate immune activation (PubMed:[28314590](#)).

Cellular Location

Cytoplasm, cytosol. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Mitochondrion Inflammasome. Secreted
Note=Predominantly localizes to the endoplasmic reticulum (ER) Association with the ER membrane requires TMEM214 (PubMed:[15123740](#)) Released in the extracellular milieu by keratinocytes following UVB irradiation (PubMed:[22246630](#)).

Tissue Location


Widely expressed, including in keratinocytes and colonic and small intestinal epithelial cells (at protein level). Not detected in brain.

Background


Inflammatory caspase (PubMed: [7797510](#), PubMed:[23516580](#), PubMed:[25119034](#)). Essential effector of NLRP3 inflammasome- dependent CASP1 activation and IL1B and IL18 secretion in response to non-canonical activators, such as UVB radiation, cholera enterotoxin subunit B and cytosolic LPS (PubMed:[22246630](#), PubMed:[26174085](#), PubMed:[26173988](#), PubMed:[26508369](#), PubMed:[25964352](#)).

Independently of NLRP3 inflammasome and CASP1, promotes pyroptosis, through GSDMD cleavage and activation, and IL1A, IL18 and HMGB1 release in response to non-canonical inflammasome activators (PubMed:[24879791](#), PubMed:[25964352](#)). Plays a crucial role in the restriction of *Salmonella typhimurium* replication in colonic epithelial cells during infection (PubMed:[25121752](#)). In later stages of the infection, LPS from cytosolic *Salmonella* triggers CASP4 activation, which ultimately results in pyroptosis of infected cells and their extrusion into the gut lumen, as well as in IL18 secretion. Pyroptosis limits bacterial replication, while cytokine secretion promotes the recruitment and activation of immune cells and triggers mucosal inflammation. Involved in LPS-induced IL6 secretion; this activity may not require caspase enzymatic activity (PubMed:[26508369](#)). Involved in cell death induced by endoplasmic reticulum stress and by treatment with cytotoxic APP peptides found Alzheimer's patient brains (PubMed:[15123740](#), PubMed:[22246630](#), PubMed:[23661706](#)). Activated by direct binding to LPS without the need of an upstream sensor (PubMed:[25119034](#)). Does not directly process IL1B (PubMed:[7743998](#), PubMed:[7797592](#), PubMed:[7797510](#)). During non- canonical inflammasome activation, cuts CGAS and may play a role in the regulation of antiviral innate immune activation (PubMed:[28314590](#)).

Images



Western Blot analysis of various cells using Cleaved-Caspase-4 p20 (Q81) Polyclonal Antibody diluted at 1 : 2000



Western Blot analysis of 3T3 cells using Cleaved-Caspase-4 p20 (Q81) Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.