

Cleaved-Cathepsin D LC (G65) Polyclonal Antibody

Catalog # AP63113

Product Information

Application	WB, E
Primary Accession	P07339
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44552

Additional Information

Gene ID	1509
Other Names	CTSD; CPSD; Cathepsin D
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	CTSD
Synonyms	CPSD
Function	Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed: 27333034). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.
Cellular Location	Lysosome. Melanosome. Secreted, extracellular space. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380)
Tissue Location	Expressed in the aorta extracellular space (at protein level) (PubMed:20551380). Expressed in liver (at protein level) (PubMed:1426530).

Background

Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed:[27333034](#)). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

Images



Western Blot analysis of various cells using
Cleaved-Cathepsin D LC (G65) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.