

# AGXT Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP6500c

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P21549</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB18848
<b>Calculated MW</b>	43010
<b>Antigen Region</b>	96-125

## Additional Information

---

<b>Gene ID</b>	189
<b>Other Names</b>	Serine--pyruvate aminotransferase, SPT, Alanine--glyoxylate aminotransferase, AGT, AGXT, AGT1, SPAT
<b>Target/Specificity</b>	This AGXT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 96-125 amino acids from the Central region of human AGXT.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	AGXT Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	AGXT ( <a href="#">HGNC:341</a> )
<b>Synonyms</b>	AGT1, SPAT
<b>Function</b>	Peroxisomal aminotransferase that catalyzes the transamination of

glyoxylate to glycine and contributes to the glyoxylate detoxification (PubMed:[10960483](#), PubMed:[12777626](#), PubMed:[23229545](#), PubMed:[24055001](#), PubMed:[26149463](#)). Also catalyzes the transamination between L-serine and pyruvate and contributes to gluconeogenesis from the L-serine metabolism (PubMed:[10347152](#)).

## Cellular Location

Peroxisome

## Tissue Location

Liver.


## Background

AGXT is expressed only in the liver and protein is localized mostly in the peroxisomes, where it is involved in glyoxylate detoxification.


## References

Cellini,B., J. Biol. Chem. 284 (13), 8349-8358 (2009)  
Bertoldi,M., Biochim. Biophys. Acta 1784 (9), 1356-1362 (2008)

## Images



AGXT Antibody (Center) (RB18848) IHC analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the AGXT Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



AGXT Antibody (Center) (Cat. #AP6500c) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.