

# CNOT8 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6521b

## Product Information

---

<b>Application</b>	IHC-P, FC, IF, WB, E
<b>Primary Accession</b>	<a href="#">Q9UFF9</a>
<b>Other Accession</b>	<a href="#">Q9D8X5</a>
<b>Reactivity</b>	Human, Mouse
<b>Predicted</b>	Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	33540
<b>Antigen Region</b>	227-255

## Additional Information

---

<b>Gene ID</b>	9337
<b>Other Names</b>	CCR4-NOT transcription complex subunit 8, CAF1-like protein, CALIFp, CAF2, CCR4-associated factor 8, Caf1b, CNOT8, CALIF, POP2
<b>Target/Specificity</b>	This CNOT8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 227-255 amino acids from the C-terminal region of human CNOT8.
<b>Dilution</b>	IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CNOT8 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CNOT8
<b>Synonyms</b>	CALIF, POP2

## Function

Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate. Its function seems to be partially redundant with that of CNOT7. Catalytic component of the CCR4-NOT complex which is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. During miRNA-mediated repression the complex also seems to act as translational repressor during translational initiation. Additional complex functions may be a consequence of its influence on mRNA expression. Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti-proliferative activity.

## Cellular Location

Cytoplasm. Nucleus

## Background


Ubiquitous transcription factor required for a diverse set of processes. The CCR4-NOT complex functions as general transcription regulation complex.

## References


Morel,A.P., J. Cell. Sci. 116 (PT 14), 2929-2936 (2003)

Prevot,D., J. Biol. Chem. 276 (13), 9640-9648 (2001)

## Images




CNOT8 Antibody (C-term) (Cat. #AP6521b) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the CNOT8 antibody detected the CNOT8 protein (arrow).




Fluorescent confocal image of HeLa cell stained with CNOT8 Antibody (C-term)(Cat#AP6521b). HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with CNOT8 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). CNOT8 immunoreactivity is localized to Cytoplasm significantly and Nucleus weakly.

Formalin-fixed and paraffin-embedded human colon carcinoma with CNOT8 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical



relevance has not been evaluated.



Flow cytometric analysis of CEM cells using CNOT8 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.