

TRAP Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6545A

Product Information

Application	WB, IHC-P, IF, FC, E
Primary Accession	P29965
Other Accession	Q95MQ5 , P51749
Reactivity	Human
Predicted	Bovine, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB19150
Calculated MW	29274
Antigen Region	33-62

Additional Information

Gene ID	959
Other Names	CD40 ligand, CD40-L, T-cell antigen Gp39, TNF-related activation protein, TRAP, Tumor necrosis factor ligand superfamily member 5, CD154, CD40 ligand, membrane form, CD40 ligand, soluble form, CD40LG, CD40L, TNFSF5, TRAP
Target/Specificity	This TRAP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 33-62 amino acids from the N-terminal region of human TRAP.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:10~50 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TRAP Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD40LG
-------------	--------

Synonyms

CD40L, TNFSF5, TRAP

Function

Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed: [1280226](#), PubMed:[31331973](#)). Costimulates T-cell proliferation and cytokine production (PubMed:[8617933](#)). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:[8617933](#)). Induces the activation of NF-kappa-B (PubMed:[15067037](#), PubMed:[31331973](#)). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:[15067037](#)). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:[15067037](#)). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell surface

Tissue Location

Specifically expressed on activated CD4+ T- lymphocytes

Background


TRAP is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in its gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome.

References


Volmar,C.H., Exp. Cell Res. 315 (13), 2265-2274 (2009)

Chai,H., Surgery 146 (1), 5-11 (2009)


Images




All lanes: Anti- TRAP Antibody (N-term) at 1:2000 dilution
Lane 1: NCI-H460 whole cell lysate Lane 2: THP-1 whole cell lysate Lane 3: HuT78 whole cell lysate Lane 4: U266B1 whole cell lysate Lysates/proteins at 20 µg per lane.
Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 29 KDa Blocking/Dilution buffer: 5% NFDm/TBST.



TRAP Antibody (N-term) (Cat.# AP6545a) IHC analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TRAP Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of TRAP Antibody (N-term)(Cat#AP6545a) with NCI-H460 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



NQO1 Antibody (Center) (Cat. #AP7350c) flow cytometric analysis of NCI-H460 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Citations

- [Circular RNA atlas in osteoclast differentiation with and without alendronate treatment](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.