

GRF-1 (phospho Tyr1105) Polyclonal Antibody

Catalog # AP67567

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9NRY4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	170514

Additional Information

Gene ID	2909
Other Names	ARHGAP35; GRF1; GRLF1; KIAA1722; Rho GTPase-activating protein 35; Glucocorticoid receptor DNA-binding factor 1; Glucocorticoid receptor repression factor 1; GRF-1; Rho GAP p190A; p190-A
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	ARHGAP35 (HGNC:4591)
Function	Rho GTPase-activating protein (GAP) (PubMed: 19673492 , PubMed: 28894085). Binds several acidic phospholipids which inhibits the Rho GAP activity to promote the Rac GAP activity (PubMed: 19673492). This binding is inhibited by phosphorylation by PRKCA (PubMed: 19673492). Involved in cell differentiation as well as cell adhesion and migration, plays an important role in retinal tissue morphogenesis, neural tube fusion, midline fusion of the cerebral hemispheres and mammary gland branching morphogenesis (By similarity). Transduces signals from p21-ras to the nucleus, acting via the ras GTPase-activating protein (GAP) (By similarity). Transduces SRC- dependent signals from cell-surface adhesion molecules, such as laminin, to promote neurite outgrowth. Regulates axon outgrowth, guidance and fasciculation (By similarity). Modulates Rho GTPase- dependent F-actin polymerization, organization and assembly, is involved in polarized cell migration and in the positive regulation of ciliogenesis and cilia elongation (By similarity). During

mammary gland development, is required in both the epithelial and stromal compartments for ductal outgrowth (By similarity). Represses transcription of the glucocorticoid receptor by binding to the cis- acting regulatory sequence 5'-GAGAAAAGAACTGGAGAACTC-3'; this function is however unclear and would need additional experimental evidences (PubMed:[1894621](#)).

Cellular Location

Cytoplasm, cytoskeleton, cilium basal body
{ECO:0000250 | UniProtKB:Q91YM2}. Cytoplasm
{ECO:0000250 | UniProtKB:Q91YM2}. Nucleus Cell membrane
{ECO:0000250 | UniProtKB:Q91YM2}. Note=In response to integrins and SDC4 and upon phosphorylation by PKC, relocates from the cytoplasm to regions of plasma membrane ruffling where it colocalizes with polymerized actin.
{ECO:0000250 | UniProtKB:Q91YM2}


Tissue Location

Detected in neutrophils (at protein level).


Background

Rho GTPase-activating protein (GAP) (PubMed: [19673492](#), PubMed: [28894085](#)). Binds several acidic phospholipids which inhibits the Rho GAP activity to promote the Rac GAP activity (PubMed: [19673492](#)). This binding is inhibited by phosphorylation by PRKCA (PubMed: [19673492](#)). Involved in cell differentiation as well as cell adhesion and migration, plays an important role in retinal tissue morphogenesis, neural tube fusion, midline fusion of the cerebral hemispheres and mammary gland branching morphogenesis (By similarity). Transduces signals from p21-ras to the nucleus, acting via the ras GTPase-activating protein (GAP) (By similarity). Transduces SRC-dependent signals from cell-surface adhesion molecules, such as laminin, to promote neurite outgrowth. Regulates axon outgrowth, guidance and fasciculation (By similarity). Modulates Rho GTPase-dependent F-actin polymerization, organization and assembly, is involved in polarized cell migration and in the positive regulation of ciliogenesis and cilia elongation (By similarity). During mammary gland development, is required in both the epithelial and stromal compartments for ductal outgrowth (By similarity). Represses transcription of the glucocorticoid receptor by binding to the cis-acting regulatory sequence 5'-GAGAAAAGAACTGGAGAACTC-3'; this function is however unclear and would need additional experimental evidences (PubMed: [1894621](#)).

Images



Western Blot analysis of 3T3 cells using Phospho-GRF-1 (Y1105) Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.