

NMDA ζ 1 (phospho Ser890) Polyclonal Antibody

Catalog # AP68067

Product Information

Application	IHC-P, IF, ICC, E
Primary Accession	Q05586
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105373

Additional Information

Gene ID	2902
Other Names	GRIN1; NMDAR1; Glutamate [NMDA] receptor subunit zeta-1; N-methyl-D-aspartate receptor subunit NR1; NMD-R1
Dilution	IHC-P~~Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	GRIN1 (HGNC:4584)
Function	Component of N-methyl-D-aspartate (NMDA) receptors (NMDARs) that function as heterotetrameric, ligand-gated cation channels with high calcium permeability and voltage-dependent block by Mg(2+) (PubMed: 21376300 , PubMed: 26875626 , PubMed: 26919761 , PubMed: 28126851 , PubMed: 28228639 , PubMed: 36959261 , PubMed: 7679115 , PubMed: 7681588 , PubMed: 7685113). NMDARs participate in synaptic plasticity for learning and memory formation by contributing to the long-term potentiation (LTP) (PubMed: 26875626). Channel activation requires binding of the neurotransmitter L-glutamate to the GluN2 subunit, glycine or D-serine binding to the GluN1 subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+) (PubMed: 21376300 , PubMed: 26875626 , PubMed: 26919761 , PubMed: 27164704 , PubMed: 28095420 , PubMed: 28105280 , PubMed: 28126851 , PubMed: 28228639 , PubMed: 36959261 , PubMed: 38538865 , PubMed: 7679115 , PubMed: 7681588 , PubMed: 7685113). NMDARs mediate simultaneously the potassium efflux and the influx of calcium and sodium (By similarity). Each GluN2 or GluN3 subunit

confers differential attributes to channel properties, including activation, deactivation and desensitization kinetics, pH sensitivity, Ca²⁺ permeability, and binding to allosteric modulators (PubMed:[26875626](#), PubMed:[26919761](#), PubMed:[36309015](#), PubMed:[38598639](#)).


Cellular Location

Cell membrane; Multi-pass membrane protein
{ECO:0000250|UniProtKB:P35439}. Postsynaptic cell membrane
{ECO:0000250|UniProtKB:P35438}. Postsynaptic density membrane
{ECO:0000250|UniProtKB:P35439}. Synaptic cell membrane
{ECO:0000250|UniProtKB:P35438}. Note=Synaptic cell membrane targeting is dependent of GRIN2B/GluN2B subunit (By similarity). Association with GRIN3A occurs in the endoplasmic reticulum (By similarity) {ECO:0000250, ECO:0000250|UniProtKB:P35438, ECO:0000250|UniProtKB:P35439}

Background

Component of NMDA receptor complexes that function as heterotetrameric, ligand-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Channel activation requires binding of the neurotransmitter glutamate to the epsilon subunit, glycine binding to the zeta subunit, plus membrane depolarization to eliminate channel inhibition by Mg²⁺ (PubMed:[7685113](#), PubMed:[28126851](#), PubMed:[26919761](#), PubMed:[26875626](#), PubMed:[28105280](#)). Sensitivity to glutamate and channel kinetics depend on the subunit composition (PubMed:[26919761](#)).

Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.