

# EpoR (phospho Tyr426) Polyclonal Antibody

Catalog # AP68137

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P19235</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	55065

## Additional Information

---

<b>Gene ID</b>	2057
<b>Other Names</b>	EPOR; Erythropoietin receptor; EPO-R
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	EPOR {ECO:0000303   PubMed:2163695, ECO:0000312   HGNC:HGNC:3416}
<b>Function</b>	Receptor for erythropoietin, which mediates erythropoietin- induced erythroblast proliferation and differentiation (PubMed: <a href="#">10388848</a> , PubMed: <a href="#">2163695</a> , PubMed: <a href="#">2163696</a> , PubMed: <a href="#">8662939</a> , PubMed: <a href="#">9774108</a> ). Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade (By similarity). In some cell types, can also activate STAT1 and STAT3 (PubMed: <a href="#">11756159</a> ). May also activate the LYN tyrosine kinase (By similarity).
<b>Cellular Location</b>	Cell membrane {ECO:0000250   UniProtKB:P14753}; Single-pass type I membrane protein
<b>Tissue Location</b>	Erythroid cells and erythroid progenitor cells. [Isoform EPOR-S]: Isoform EPOR-S and isoform EPOR-T are the predominant forms in bone marrow.

## Background


---

Receptor for erythropoietin. Mediates erythropoietin- induced erythroblast proliferation and differentiation. Upon EPO stimulation, EPOR dimerizes triggering the JAK2/STAT5 signaling cascade. In some

cell types, can also activate STAT1 and STAT3. May also activate the LYN tyrosine kinase.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.