

C1INH Polyclonal Antibody

Catalog # AP68744

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P05155
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55154

Additional Information

Gene ID	710
Other Names	SERPING1; C1IN; C1NH; Plasma protease C1 inhibitor; C1 Inh; C1Inh; C1 esterase inhibitor; C1-inhibiting factor; Serpin G1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	SERPING1
Synonyms	C1IN, C1NH
Function	Serine protease inhibitor, which acts as a regulator of the classical complement pathway (PubMed: 10946292 , PubMed: 11527969 , PubMed: 3458172 , PubMed: 6416294). Forms a proteolytically inactive stoichiometric complex with the C1r or C1s proteases (PubMed: 10946292 , PubMed: 3458172 , PubMed: 6416294). May also regulate blood coagulation, fibrinolysis and the generation of kinins (PubMed: 8495195). Very efficient inhibitor of FXIIa. Inhibits chymotrypsin and kallikrein (PubMed: 8495195).
Cellular Location	Secreted

Background


Activation of the C1 complex is under control of the C1- inhibitor. It forms a proteolytically inactive

stoichiometric complex with the C1r or C1s proteases. May play a potentially crucial role in regulating important physiological pathways including complement activation, blood coagulation, fibrinolysis and the generation of kinins. Very efficient inhibitor of FXIIa. Inhibits chymotrypsin and kallikrein.

Images



Western Blot analysis of various cells using C1INH Polyclonal Antibody



Western Blot analysis of HuvEc cells using C1INH Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.