

CA II Polyclonal Antibody

Catalog # AP68756

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P00918
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29246

Additional Information

Gene ID	760
Other Names	CA2; Carbonic anhydrase 2; Carbonate dehydratase II; Carbonic anhydrase C; CAC; Carbonic anhydrase II; CA-II
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CA2
Function	Catalyzes the reversible hydration of carbon dioxide (PubMed: 11327835 , PubMed: 11802772 , PubMed: 11831900 , PubMed: 12056894 , PubMed: 12171926 , PubMed: 1336460 , PubMed: 14736236 , PubMed: 15300855 , PubMed: 15453828 , PubMed: 15667203 , PubMed: 15865431 , PubMed: 16106378 , PubMed: 16214338 , PubMed: 16290146 , PubMed: 16686544 , PubMed: 16759856 , PubMed: 16807956 , PubMed: 17127057 , PubMed: 17251017 , PubMed: 17314045 , PubMed: 17330962 , PubMed: 17346964 , PubMed: 17540563 , PubMed: 17588751 , PubMed: 17705204 , PubMed: 18024029 , PubMed: 18162396 , PubMed: 18266323 , PubMed: 18374572 , PubMed: 18481843 , PubMed: 18618712 , PubMed: 18640037 , PubMed: 18942852 , PubMed: 1909891 , PubMed: 1910042 , PubMed: 19170619 , PubMed: 19186056 , PubMed: 19206230 , PubMed: 19520834 , PubMed: 19778001 , PubMed: 7761440 , PubMed: 7901850 , PubMed: 8218160 , PubMed: 8262987 , PubMed: 8399159 , PubMed: 8451242 , PubMed: 8485129 , PubMed: 8639494 , PubMed: 9265618 , PubMed: 9398308). Can also hydrate

cyanamide to urea (PubMed:[10550681](#), PubMed:[11015219](#)). Stimulates the chloride-bicarbonate exchange activity of SLC26A6 (PubMed:[15990874](#)). Essential for bone resorption and osteoclast differentiation (PubMed:[15300855](#)). Involved in the regulation of fluid secretion into the anterior chamber of the eye. Contributes to intracellular pH regulation in the duodenal upper villous epithelium during proton-coupled peptide absorption.


Cellular Location

Cytoplasm. Cell membrane. Note=Colocalized with SLC26A6 at the surface of the cell membrane in order to form a bicarbonate transport metabolon. Displaced from the cytosolic surface of the cell membrane by PKC in phorbol myristate acetate (PMA)-induced cells

Background

Essential for bone resorption and osteoclast differentiation (By similarity). Reversible hydration of carbon dioxide. Can hydrate cyanamide to urea. Involved in the regulation of fluid secretion into the anterior chamber of the eye. Contributes to intracellular pH regulation in the duodenal upper villous epithelium during proton-coupled peptide absorption. Stimulates the chloride-bicarbonate exchange activity of SLC26A6.

Images



Western blot analysis of Mouse-kidney mouse-brain 293T hela lysis using CA II antibody. Antibody was diluted at 1:2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.