

Casein Kinase Iε Polyclonal Antibody

Catalog # AP68834

Product Information

Application	WB, IHC-P, IF
Primary Accession	P49674
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47315

Additional Information

Gene ID	1454
Other Names	CSNK1E; Casein kinase I isoform epsilon; CKI-epsilon; CKIε
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	CSNK1E
Function	Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates (Probable). Participates in Wnt signaling (PubMed: 12556519 , PubMed: 23413191). Phosphorylates DVL1 (PubMed: 12556519). Phosphorylates DVL2 (PubMed: 23413191). Phosphorylates NEDD9/HEF1 (By similarity). Central component of the circadian clock (PubMed: 16790549). In balance with PP1, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation (PubMed: 15917222 , PubMed: 16790549). Controls PER1 and PER2 nuclear transport and degradation (By similarity). Inhibits cytokine-induced granulocytic differentiation (PubMed: 15070676).
Cellular Location	Cytoplasm. Nucleus.
Tissue Location	Expressed in all tissues examined, including brain, heart, lung, liver, pancreas, kidney, placenta and skeletal muscle Expressed in monocytes and lymphocytes but not in granulocytes

Background

Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. Can phosphorylate a large number of proteins. Participates in Wnt signaling. Phosphorylates DVL1 and DVL2. Central component of the circadian clock. In balance with PP1, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. Controls PER1 and PER2 nuclear transport and degradation. Inhibits cytokine-induced granulocytic differentiation.

Images



Western Blot analysis of LOVO cells using Casein Kinase Iε Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.