

CIDE-B Polyclonal Antibody

Catalog # AP69111

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9UHD4
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24678

Additional Information

Gene ID	27141
Other Names	CIDEB; Cell death activator CIDE-B; Cell death-inducing DFFA-like effector B
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50-200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CIDEB {ECO:0000303 PubMed:35939579, ECO:0000312 HGNC:HGNC:1977}
Function	Lipid transferase specifically expressed in hepatocytes, which promotes unilocular lipid droplet formation by mediating lipid droplet fusion (PubMed: 35939579). Lipid droplet fusion promotes their enlargement, restricting lipolysis and favoring lipid storage (PubMed: 35939579). Localizes on the lipid droplet surface, at focal contact sites between lipid droplets, and mediates atypical lipid droplet fusion by promoting directional net neutral lipid transfer from the smaller to larger lipid droplets (By similarity). The transfer direction may be driven by the internal pressure difference between the contacting lipid droplet pair (By similarity). Promotes lipid exchange and lipid droplet fusion in both small and large lipid droplet- containing hepatocytes (By similarity). In addition to its role in lipid droplet fusion, also involved in cytoplasmic vesicle biogenesis and transport (By similarity). Required for very-low-density lipoprotein (VLDL) lipidation and maturation (By similarity). Probably involved in the biogenesis of VLDL transport vesicles by forming a COPII vesicle coat and facilitating the formation of endoplasmic reticulum-derived large vesicles (By similarity). Also involved in sterol-regulated export of the SCAP-SREBP complex, composed of SCAP,

SREBF1/SREBP1 and SREBF2/SREBP2, by promoting loading of SCAP-SREBP into COPII vesicles (By similarity). May also activate apoptosis (PubMed:[10619428](#)).

Cellular Location

Lipid droplet. Endoplasmic reticulum membrane
{ECO:0000250|UniProtKB:O70303}; Peripheral membrane protein
{ECO:0000250|UniProtKB:O70303}; Cytoplasmic side
{ECO:0000250|UniProtKB:O70303}. Golgi apparatus
{ECO:0000250|UniProtKB:O70303}. Cytoplasmic vesicle, COPII-coated vesicle
{ECO:0000250|UniProtKB:O70303}. Note=Enriched at lipid droplet contact sites. {ECO:0000250|UniProtKB:O70303}


Tissue Location

Highly expressed in liver and small intestine and, at lower levels, in colon, kidney and spleen

Background

Activates apoptosis.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.