

# Cyclin D3 Polyclonal Antibody

Catalog # AP69358

## Product Information

---

|                          |                           |
|--------------------------|---------------------------|
| <b>Application</b>       | WB, IHC-P, IF, ICC, E     |
| <b>Primary Accession</b> | <a href="#">P30281</a>    |
| <b>Reactivity</b>        | Human, Mouse, Rat, Monkey |
| <b>Host</b>              | Rabbit                    |
| <b>Clonality</b>         | Polyclonal                |
| <b>Calculated MW</b>     | 32520                     |

## Additional Information

---

|                           |   |
|---------------------------|---|
| <b>Gene ID</b>            | 896   |
| <b>Other Names</b>        | CCND3; G1/S-specific cyclin-D3  |
| <b>Dilution</b>           | WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A |
| <b>Format</b>             | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.   |
| <b>Storage Conditions</b> | -20°C   |

## Protein Information


---

|                          |   |
|--------------------------|---|
| <b>Name</b>              | CCND3 {ECO:0000303   PubMed:1386336, ECO:0000312   HGNC:HGNC:1585}  |
| <b>Function</b>          | Regulatory component of the cyclin D3-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: <a href="#">8114739</a> ). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed: <a href="#">8114739</a> ). Hypophosphorylates RB1 in early G(1) phase (PubMed: <a href="#">8114739</a> ). Cyclin D- CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed: <a href="#">8114739</a> ). Component of the ternary complex, cyclin D3/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (PubMed: <a href="#">16782892</a> ). Shows transcriptional coactivator activity with ATF5 independently of CDK4 (PubMed: <a href="#">15358120</a> ). |
| <b>Cellular Location</b> | Nucleus. Cytoplasm  |


## Background

Regulatory component of the cyclin D3-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D3/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex.

## Images



Western Blot analysis of COS7 cells using Cyclin D3 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human tonsil. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.