

# CYP2S1 Polyclonal Antibody

Catalog # AP69406

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q96SQ9</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	55817

## Additional Information

---

<b>Gene ID</b>	29785
<b>Other Names</b>	CYP2S1; Cytochrome P450 2S1; CYPIIS1
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50-200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	CYP2S1 {ECO:0000303   PubMed:11181079, ECO:0000312   HGNC:HGNC:15654}
<b>Function</b>	A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed: <a href="#">12711469</a> , PubMed: <a href="#">21068195</a> ). In epidermis, may contribute to the oxidative metabolism of all-trans- retinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed: <a href="#">12711469</a> ). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates (HPETEs) (PubMed: <a href="#">21068195</a> ). Independently of cytochrome P450 reductase, NADPH, and O2, catalyzes the breakdown of PGH2 to hydroxyheptadecatrienoic acid (HHT) and malondialdehyde (MDA), which is known to act as a mediator of DNA damage (PubMed: <a href="#">21068195</a> ).
<b>Cellular Location</b>	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein


## Tissue Location

Expressed at higher levels in extrahepatic tissues including trachea, lung, stomach, small intestine, colon, kidney, breast, placenta and spleen (PubMed:11181079, PubMed:12711469) Expressed in peripheral blood leukocytes (PubMed:11181079) Constitutively expressed in skin (at protein level) (PubMed:12711469)

## Background

Has a potential importance for extrahepatic xenobiotic metabolism.

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.