

# CYP2W1 Polyclonal Antibody

Catalog # AP69408

## Product Information

Application	WB, IHC-P, IF
Primary Accession	<a href="#">Q8TAV3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53844

## Additional Information

Gene ID	54905
Other Names	CYP2W1; Cytochrome P450 2W1; CYP11W1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

Name	CYP2W1 {ECO:0000303   PubMed:26936974, ECO:0000312   HGNC:HGNC:20243}
Function	<p>A cytochrome P450 monooxygenase that may play a role in retinoid and phospholipid metabolism (PubMed:<a href="#">22591743</a>, PubMed:<a href="#">26936974</a>). Catalyzes the hydroxylation of saturated carbon hydrogen bonds. Hydroxylates all trans-retinoic acid (atRA) to 4- hydroxyretinoate and may regulate atRA clearance. Other retinoids such as all-trans retinol and all-trans retinal are potential endogenous substrates (PubMed:<a href="#">26936974</a>). Catalyzes both epoxidation of double bonds and hydroxylation of carbon hydrogen bonds of the fatty acyl chain of 1-acylphospholipids/2-lysophospholipids. Can metabolize various lysophospholipids classes including lysophosphatidylcholines (LPCs), lysophosphatidylinositols (LPIs), lysophosphatidylserines (LPSs), lysophosphatidylglycerols (LPGs), lysophosphatidylethanolamines (LPEs) and lysophosphatidic acids (LPAs) (PubMed:<a href="#">22591743</a>). Has low or no activity toward 2-acylphospholipids/1-lysophospholipids, diacylphospholipids and free fatty acids (PubMed:<a href="#">22591743</a>, PubMed:<a href="#">26936974</a>). May play a role in tumorigenesis by activating procarcinogens such as aflatoxin B1, polycyclic</p>

aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:[16551781](#), PubMed:[20805301](#), PubMed:[24278521](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:[22591743](#), PubMed:[26936974](#)).

**Cellular Location**

Endoplasmic reticulum lumen. Cell membrane. Microsome membrane.  
Note=About 8% are expressed on the cell surface.

**Tissue Location**

Very low levels are detected in fetal and adult tissues. Highly expressed in several tumor samples, in particular colon and adrenal tumors.


## Background

---

Seems to have broad catalytic activity towards several chemicals, including polycyclic aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:[16551781](#), PubMed:[24278521](#)). Active also in the metabolism of indoline substrates and is able to activate aflatoxin B1 into cytotoxic products (PubMed:[20805301](#)). Furthermore, it seems to be involved in the oxydation of lysophospholipids and fatty acids (PubMed:[22591743](#)).

## Images

---



Western Blot analysis of various cells using CYP2W1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.