

FN1 Polyclonal Antibody

Catalog # AP69916

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P02751
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	272320

Additional Information

Gene ID	2335
Other Names	FN1; FN; Fibronectin; FN; Cold-insoluble globulin; CIG
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	FN1 (HGNC:3778)
Synonyms	FN
Function	Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed: 3024962 , PubMed: 3593230 , PubMed: 3900070 , PubMed: 7989369). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed: 3024962 , PubMed: 3593230 , PubMed: 3900070 , PubMed: 7989369). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor, inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed: 34089617).
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted {ECO:0000250 UniProtKB:P11276}


Tissue Location

Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).


Background

Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin. Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape. Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization. Participates in the regulation of type I collagen deposition by osteoblasts.


Images



Western Blot analysis of various cells using FN1 Polyclonal Antibody diluted at 1 : 2000



Western Blot analysis of HeLa cells using FN1 Polyclonal Antibody diluted at 1 : 2000



Western blot analysis of 3T3 KB K562 293T lysis using FN1 antibody. Antibody was diluted at 1:2000