

# Follistatin Polyclonal Antibody

Catalog # AP69918

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">P19883</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	38007

## Additional Information

---

<b>Gene ID</b>	10468
<b>Other Names</b>	FST; Follistatin; FS; Activin-binding protein
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	FST ( <a href="#">HGNC:3971</a> )
<b>Function</b>	Multifunctional regulatory protein whose primary function is to antagonize members of the transforming growth factor beta (TGF-beta) superfamily including activin, myostatin, GDF11 or bone morphogenetic proteins (BMPs) (PubMed: <a href="#">11279126</a> , PubMed: <a href="#">16482217</a> , PubMed: <a href="#">18535106</a> ). Mechanistically, binds to these ligands in the extracellular space, blocking their type II receptor-binding site to inhibit downstream signaling (PubMed: <a href="#">16482217</a> ). Plays an essential role in muscle fiber formation and growth both by preventing the repressive effects of myostatin and through SMAD3/AKT/mTOR signaling independently of myostatin (By similarity). Also promotes neural differentiation by antagonizing the action BMP4 (By similarity). Acts as a specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone (FSH) by sequestering activin A/INHBA (PubMed: <a href="#">11279126</a> ). On the other hand, translocates into the nucleus where it down-regulates rRNA synthesis and ribosome biogenesis to maintain cellular energy homeostasis by binding to rDNA.
<b>Cellular Location</b>	Secreted. Nucleus, nucleolus

## Tissue Location

Isoform 1 is the predominant isoform in serum but is undetectable in follicular fluid. In the embryo, strong expression is seen in the palatal epithelia, including the medial edge epithelial and midline epithelial seam of the palatal shelves. Less pronounced expression is also seen throughout the palatal shelf and tongue mesenchyme (PubMed:31215115).


## Background

---

Binds directly to activin and functions as an activin antagonist. Specific inhibitor of the biosynthesis and secretion of pituitary follicle stimulating hormone (FSH).

## Images

---



Western Blot analysis of various cells using Follistatin Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.