

FoxN4 Polyclonal Antibody

Catalog # AP69939

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q96NZ1
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55215

Additional Information

Gene ID	121643
Other Names	FOXN4; Forkhead box protein N4
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	FOXN4
Function	Transcription factor essential for neural and some non-neural tissues development, such as retina and lung respectively. Binds to an 11-bp consensus sequence containing the invariant tetranucleotide 5'- ACGC-3'. During development of the central nervous system, is required to specify the amacrine and horizontal cell fates from multipotent retinal progenitors while suppressing the alternative photoreceptor cell fates through activating DLL4-NOTCH signaling. Also acts synergistically with ASCL1/MASH1 to activate DLL4-NOTCH signaling and drive commitment of p2 progenitors to the V2b interneuron fates during spinal cord neurogenesis. In development of non-neural tissues, plays an essential role in the specification of the atrioventricular canal and is indirectly required for patterning the distal airway during lung development (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00089}.

Background

Transcription factor essential for neural and some non- neural tissues development, such as retina and lung respectively. Binds to an 11-bp consensus sequence containing the invariant tetranucleotide 5'-ACGC-3'. During development of the central nervous system, is required to specify the amacrine and horizontal cell fates from multipotent retinal progenitors while suppressing the alternative photoreceptor cell fates through activating DLL4- NOTCH signaling. Also acts synergistically with ASCL1/MASH1 to activate DLL4-NOTCH signaling and drive commitment of p2 progenitors to the V2b interneuron fates during spinal cord neurogenesis. In development of non-neural tissues, plays an essential role in the specification of the atrioventricular canal and is indirectly required for patterning the distal airway during lung development (By similarity).

Images



Western Blot analysis of LOVO cells using FoxN4 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.