

GPR18 Polyclonal Antibody

Catalog # AP70187

Product Information

Application	WB, IF
Primary Accession	Q14330
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38134

Additional Information

Gene ID	2841
Other Names	GPR18; GPCRW; N-arachidonyl glycine receptor; NAGly receptor; G-protein coupled receptor 18
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	GPR18
Synonyms	GPCRW
Function	<p>G protein-coupled receptor (GPCR) that plays a role in diverse physiological processes particularly within the immune and nervous systems (PubMed:21732409, PubMed:26195725). Becomes active when triggered by various endogenous ligands including endocannabinoid N- arachidonyl glycine (NAGly), delta-9-tetrahydrocannabinol or resolvin D2/RvD2 derived from the omega-3 fatty acid docosahexaenoic acid (DHA) (PubMed:16844083, PubMed:24762058, PubMed:26195725, PubMed:27572937). Upon RvD2 binding, facilitates the resolution of inflammation, aiding in tissue repair and homeostasis. Mechanistically, RvD2 ligation initiates Gα_i protein coupling, activation of cAMP-PKA signaling pathway and phosphorylation of STAT3, leading to RvD2-stimulated macrophage phagocytosis (PubMed:27994074). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937). Activation by N-arachidonoyl glycine (NAGly) can also induce apoptosis in macrophages (By similarity). Plays a role in homeostasis of CD8+ subsets of intraepithelial</p>

lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs) in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Also participates in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane

Tissue Location

Expressed in midpiece of spermatozoon (at protein level) (PubMed:27572937). Most abundant in testis and spleen (PubMed:16844083). Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells (PubMed:16844083)

Background

Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (PubMed: [16844083](#), PubMed:[24762058](#), PubMed:[27572937](#)). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:[27018161](#)). Can also be activated by plant- derived and synthetic cannabinoid agonists (PubMed:[24762058](#)). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:[16844083](#)). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8+ subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:[27572937](#)).

Images



Western Blot analysis of various cells using GPR18 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.