

Ksr-1 Polyclonal Antibody

Catalog # AP70679

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q8IVT5
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102160

Additional Information

Gene ID	8844
Other Names	KSR1; KSR; Kinase suppressor of Ras 1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50-200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	KSR1
Synonyms	KSR
Function	Part of a multiprotein signaling complex which promotes phosphorylation of Raf family members and activation of downstream MAP kinases (By similarity). Independently of its kinase activity, acts as MAP2K1/MEK1 and MAP2K2/MEK2-dependent allosteric activator of BRAF; upon binding to MAP2K1/MEK1 or MAP2K2/MEK2, dimerizes with BRAF and promotes BRAF-mediated phosphorylation of MAP2K1/MEK1 and/or MAP2K2/MEK2 (PubMed: 29433126). Promotes activation of MAPK1 and/or MAPK3, both in response to EGF and to cAMP (By similarity). Its kinase activity is unsure (By similarity). Some protein kinase activity has been detected in vitro, however the physiological relevance of this activity is unknown (By similarity).
Cellular Location	Cytoplasm. Membrane; Peripheral membrane protein. Cell membrane {ECO:0000250 UniProtKB:Q61097}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q61097}. Cell projection, ruffle membrane {ECO:0000250 UniProtKB:Q61097}. Endoplasmic reticulum membrane.

Note=In unstimulated cells, where the phosphorylated form is bound to a 14-3-3 protein, sequestration in the cytoplasm occurs. Following growth factor treatment, the protein is free for membrane translocation, and it moves from the cytoplasm to the cell periphery.

Background

Scaffolding protein that is part of a multiprotein signaling complex. Promotes phosphorylation of Raf family members and activation of downstream MAP kinases. Promotes activation of MAPK1 and/or MAPK3, both in response to EGF and to cAMP. Does not have kinase activity by itself.

Images



Western Blot analysis of NIH-3T3 cells using Ksr-1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.