

MC2-R Polyclonal Antibody

Catalog # AP70857

Product Information

Application	WB, E, IHC-P
Primary Accession	Q01718
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33927

Additional Information

Gene ID	4158
Other Names	MC2R; ACTHR; Adrenocorticotropin hormone receptor; ACTH receptor; ACTH-R; Adrenocorticotropin receptor; Melanocortin receptor 2; MC2-R
Dilution	WB--Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. E--N/A IHC-P--N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	MC2R (HGNC:6930)
Synonyms	ACTHR
Function	G protein-coupled receptor for corticotropin/ACTH, primarily expressed in adrenal cortex where it plays a key role in the regulation of adrenocortical function (PubMed: 1325670 , PubMed: 17596328 , PubMed: 36588120). Upon activation, couples to G(s) protein, stimulating adenylate cyclase and activating the cAMP-dependent signaling pathway, the MAPK cascade as well as the PKA pathway, leading to steroidogenic factor 1/NR5A1-mediated transcriptional activation (PubMed: 1325670). Activation by ACTH facilitates the release of adrenal glucocorticoids, including cortisol and corticosterone (PubMed: 12213892 , PubMed: 8636348). In addition, MC2R is required for fetal and neonatal adrenal gland development (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Melanocytes and corticoadrenal tissue.

Background

Receptor for corticotropin (ACTH). This receptor is mediated by G proteins (G(s)) which activate adenylate cyclase (cAMP).

Images



Western Blot analysis of various cells using MC2-R Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.