

MEK-3 Polyclonal Antibody

Catalog # AP70900

Product Information

Application	WB, E, IHC-P
Primary Accession	P46734
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39318

Additional Information

Gene ID	5606
Other Names	MAP2K3; MEK3; MKK3; PRKMK3; SKK2; Dual specificity mitogen-activated protein kinase kinase 3; MAP kinase kinase 3; MAPKK 3; MAPK/ERK kinase 3; MEK 3; Stress-activated protein kinase kinase 2; SAPK kinase 2; SAPKK-2; SAPKK2
Dilution	WB--Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. E--N/A IHC-P--N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	MAP2K3
Synonyms	MEK3, MKK3, PRKMK3, SKK2
Function	Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14.
Tissue Location	Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues

Background


Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the

concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14.

Images



Western Blot analysis of various cells using MEK-3 Polyclonal Antibody



Western Blot analysis of K562 cells using MEK-3 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.