

Neuralized-1 Polyclonal Antibody

Catalog # AP71235

Product Information

Application	WB, IHC-P
Primary Accession	O76050
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61860

Additional Information

Gene ID	9148
Other Names	NEURL; NEURL1; NEURL1A; RNF67; Neuralized-like protein 1A; h-neu; h-neuralized 1; RING finger protein 67
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	NEURL1
Synonyms	NEURL, NEURL1A, RNF67
Function	Plays a role in hippocampal-dependent synaptic plasticity, learning and memory. Involved in the formation of spines and functional synaptic contacts by modulating the translational activity of the cytoplasmic polyadenylation element-binding protein CPEB3. Promotes ubiquitination of CPEB3, and hence induces CPEB3-dependent mRNA translation activation of glutamate receptor GRIA1 and GRIA2. Can function as an E3 ubiquitin-protein ligase to activate monoubiquitination of JAG1 (in vitro), thereby regulating the Notch pathway. Acts as a tumor suppressor; inhibits malignant cell transformation of medulloblastoma (MB) cells by inhibiting the Notch signaling pathway.
Cellular Location	Cytoplasm, perinuclear region. Cell membrane; Peripheral membrane protein Perikaryon. Cell projection, dendrite Postsynaptic density. Note=Localized in the cell bodies of the pyramidal neurons and distributed along their apical dendrites Colocalized with PSD95 in postsynaptic sites. Colocalized with CPEB3 at apical dendrites of CA1 neurons (By similarity). Colocalized with

JAG1 at the cell surface.


Tissue Location

Expressed in brain, testis, pituitary gland, pancreas and bone marrow. Also poorly expressed in malignant astrocytomas and several neuroectodermal tumor cell lines. Weakly expressed in medulloblastoma (MB) compared with normal cerebellar tissues.

Background

Plays a role in hippocampal-dependent synaptic plasticity, learning and memory. Involved in the formation of spines and functional synaptic contacts by modulating the translational activity of the cytoplasmic polyadenylation element- binding protein CPEB3. Promotes ubiquitination of CPEB3, and hence induces CPEB3-dependent mRNA translation activation of glutamate receptor GRIA1 and GRIA2. Can function as an E3 ubiquitin-protein ligase to activate monoubiquitination of JAG1 (in vitro), thereby regulating the Notch pathway. Acts as a tumor suppressor; inhibits malignant cell transformation of medulloblastoma (MB) cells by inhibiting the Notch signaling pathway.

Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.