

PGK1 Antibody (Center S320)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7169a

Product Information

Application	IHC-P, WB, E
Primary Accession	P00558
Other Accession	Q60HD8 , Q5J7W1
Reactivity	Human, Rat, Mouse
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	44615
Antigen Region	305-334

Additional Information

Gene ID	5230
Other Names	Phosphoglycerate kinase 1, Cell migration-inducing gene 10 protein, Primer recognition protein 2, PRP 2, PGK1, PGKA
Target/Specificity	This PGK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 305-334 amino acids from the Central region of human PGK1.
Dilution	IHC-P~1:100~500 WB~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PGK1 Antibody (Center S320) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PGK1 (HGNC:8896)
Synonyms	PGKA
Function	Catalyzes one of the two ATP producing reactions in the glycolytic pathway

via the reversible conversion of 1,3- diphosphoglycerate to 3-phosphoglycerate (PubMed:[30323285](#), PubMed:[7391028](#)). Both L- and D- forms of purine and pyrimidine nucleotides can be used as substrates, but the activity is much lower on pyrimidines (PubMed:[18463139](#)). In addition to its role as a glycolytic enzyme, it seems that PGK1 acts as a polymerase alpha cofactor protein (primer recognition protein) (PubMed:[2324090](#)). Acts as a protein kinase when localized to the mitochondrion where it phosphorylates pyruvate dehydrogenase kinase PDK1 to inhibit pyruvate dehydrogenase complex activity and suppress the formation of acetyl- coenzyme A from pyruvate, and consequently inhibit oxidative phosphorylation and promote glycolysis (PubMed:[26942675](#), PubMed:[36849569](#)). May play a role in sperm motility (PubMed:[26677959](#)).

Cellular Location

Cytoplasm, cytosol. Mitochondrion matrix. Note=Hypoxic conditions promote mitochondrial targeting (PubMed:26942675). Targeted to the mitochondrion following phosphorylation by MAPK1/ERK2, cis-trans isomerization by PIN1, and binding to mitochondrial circRNA mcPGK1 (PubMed:36849569).

Tissue Location

Mainly expressed in spermatogonia. Localized on the principle piece in the sperm (at protein level). Expression significantly decreased in the testis of elderly men


Background

Also known as ATP:3-phosphoglycerate 1-phosphotransferase, this major enzyme in glycolysis catalyzes the reversible conversion of 1,3-diphosphoglycerate to 3-phosphoglycerate, generating one molecule of ATP. Phosphoglycerate kinase not only functions in glycolysis but is secreted by tumor cells and is proposed to participate in the angiogenic process as a disulfide reductase. Mutations in PGK1 may be associated with hemolytic anemia.

References


Shetty, S., et al., Am. J. Respir. Cell Mol. Biol. 31(1):100-106 (2004).
Daly, E.B., et al., Biochim. Biophys. Acta 1691(1):17-22 (2004).
Daly, E.B., et al., Int. J. Biol. Markers 19(2):170-172 (2004).
Saito, Y., et al., Biochem. Biophys. Res. Commun. 314(2):396-402 (2004).
Krishnan, P., et al., J. Biol. Chem. 278(38):36726-36732 (2003).

Images



Western blot analysis of lysates from A431, HeLa, U-937, BA/F3 cell line (from left to right), using PGK1 Antibody (S320) (Cat. #AP7169a). AP7169a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with PGK1 Antibody (Center S320)(Cat.#AP7169a), which was



peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Citations

- [A biotinylated analog of the anti-proliferative prostaglandin A1 allows assessment of PPAR-independent effects and identification of novel cellular targets for covalent modification.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.