

# ELF4 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7185c

## Product Information

---

Application	FC, WB, E
Primary Accession	<a href="#">Q99607</a>
Other Accession	<a href="#">Q9Z2U4</a> , <a href="#">Q60775</a> , <a href="#">P32519</a> , <a href="#">A0JN51</a>
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	70730
Antigen Region	211-237

## Additional Information

---

Gene ID	2000
Other Names	ETS-related transcription factor Elf-4, E74-like factor 4, Myeloid Elf-1-like factor, ELF4 {ECO:0000312   EMBL:CAI428821}
Target/Specificity	This ELF4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 211-237 amino acids from the Central region of human ELF4.
Dilution	FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ELF4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	ELF4 {ECO:0000312   EMBL:CAI42882.1}
Function	Transcriptional activator that binds to DNA sequences containing the consensus 5'-WGGA-3'. Transactivates promoters of the hematopoietic growth factor genes CSF2, IL3, IL8, and of the bovine lysozyme gene. Acts

synergistically with RUNX1 to transactivate the IL3 promoter (By similarity). Transactivates the PRF1 promoter in natural killer (NK) cells and CD8+ T cells (PubMed:[34326534](#)). Plays a role in the development and function of NK and NK T-cells and in innate immunity. Controls the proliferation and homing of CD8+ T-cells via the Kruppel-like factors KLF4 and KLF2 (By similarity). Controls cell senescence in a p53-dependent manner. Can also promote cellular transformation through inhibition of the p16 pathway. Is a transcriptional regulator of inflammation, controlling T-helper 17 (Th17) cells and macrophage inflammatory responses. Required for sustained transcription of anti-inflammatory genes, including IL1RN (PubMed:[34326534](#), PubMed:[35266071](#)). Is a negative regulator of pro- inflammatory cytokines expression including IL17A, IL1B, IL6, TNFA and CXCL1 (PubMed:[34326534](#), PubMed:[35266071](#)). Down-regulates expression of TREM1, a cell surface receptor involved in the amplification of inflammatory responses (By similarity) (PubMed:[34326534](#), PubMed:[35266071](#)).

#### Cellular Location

Nucleus, PML body. Note=Accumulation into PML nuclear bodies is mediated by PML

#### Tissue Location

Abundantly expressed in the placenta and in a variety of myeloid leukemia cell lines. Moderate levels of expression in heart, lung, spleen, thymus, peripheral blood lymphocytes, ovary and colon. Lower levels of expression in Jurkat T-cells and other T-cell lines and no expression in brain.


## Background

ELF4 is transcriptional activator that binds to DNA sequences containing the consensus 5'-WGGA-3'. The protein acts synergistically with RUNX1 to transactivate the IL3 promoter. Also transactivates the PRF1 promoter in natural killer (NK) cells. This protein plays a role in the development and function of NK and NK T-cells and in innate immunity.

## References


Miyazaki Y., Sun X.Oncogene 13:1721-1729(1996)  
 Aryee D.N.T., Petermann R.Gene 210:71-78(1998)  
 Mao S., Frank R.C., Zhang J.Mol. Cell. Biol. 19:3635-3644(1999)

## Images



ELF4 Antibody (Center) (Cat. #AP7185c) western blot analysis in HL-60,K562 cell line lysates (35ug/lane).This demonstrates the ELF4 antibody detected the ELF4 protein (arrow).

Flow cytometric analysis of NCI-H292 cells using ELF4 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the



analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.