

PFK-C Polyclonal Antibody

Catalog # AP71869

Product Information

Application WB, IHC-P, IF **Primary Accession** Q01813

Reactivity Human, Mouse, Rat, Monkey

HostRabbitClonalityPolyclonalCalculated MW85596

Additional Information

Gene ID 5214

Other Names PFKP; PFKF; 6-phosphofructokinase type C; 6-phosphofructokinase; platelet

type; Phosphofructo-1-kinase isozyme C; PFK-C; Phosphofructokinase 1;

Phosphohexokinase

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet

tested in other applications. IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name PFKP

Synonyms PFKF

Function Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose


1,6-bisphosphate by ATP, the first committing step of glycolysis.

Cellular Location Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03184}.


Background

Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose 1,6-bisphosphate by ATP, the first committing step of glycolysis.


Images



Western Blot analysis of various cells using PFK-C Polyclonal Antibody



Western Blot analysis of HuvEc cells using PFK-C Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human testis. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.