

PJA2 Polyclonal Antibody

Catalog # AP71925

Product Information

Application	WB, E, IHC-P
Primary Accession	O43164
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78214

Additional Information

Gene ID	9867
Other Names	PJA2; KIAA0438; RNF131; E3 ubiquitin-protein ligase Praja-2; Praja2; RING finger protein 131
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. E~~N/A IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information



Name	PJA2
Synonyms	KIAA0438, RNF131
Function	Has E2-dependent E3 ubiquitin-protein ligase activity (PubMed: 12036302 , PubMed: 21423175). Responsible for ubiquitination of cAMP-dependent protein kinase type I and type II-alpha/beta regulatory subunits and for targeting them for proteasomal degradation. Essential for PKA-mediated long-term memory processes (PubMed: 21423175). Through the ubiquitination of MFHAS1, positively regulates the TLR2 signaling pathway that leads to the activation of the downstream p38 and JNK MAP kinases and promotes the polarization of macrophages toward the pro- inflammatory M1 phenotype (PubMed: 28471450). Plays a role in ciliogenesis by ubiquitinating OFD1 (PubMed: 33934390).
Cellular Location	Cytoplasm. Cell membrane. Endoplasmic reticulum membrane; Peripheral membrane protein. Golgi apparatus membrane; Peripheral membrane protein. Synapse {ECO:0000250 UniProtKB:Q63364} Postsynaptic density {ECO:0000250 UniProtKB:Q63364}. Cytoplasm, cytoskeleton, microtubule

organizing center, centrosome. Note=Localizes at the cytoplasmic side of endoplasmic reticulum and Golgi apparatus (PubMed:21423175) Expressed in the postsynaptic density region of synapses (By similarity). Colocalizes with PRKAR2A and PRKAR2B in the cytoplasm and the cell membrane (PubMed:21423175). {ECO:0000250 | UniProtKB:Q63364, ECO:0000269 | PubMed:21423175}

Background

Has E2-dependent E3 ubiquitin-protein ligase activity. Responsible for ubiquitination of cAMP-dependent protein kinase type I and type II-alpha/beta regulatory subunits and for targeting them for proteasomal degradation. Essential for PKA- mediated long-term memory processes. Through the ubiquitination of MFHAS1, positively regulates the TLR2 signaling pathway that leads to the activation of the downstream p38 and JNK MAP kinases and promotes the polarization of macrophages toward the pro- inflammatory M1 phenotype (PubMed:[28471450](#)).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.