

# PTH/PTHrP-R Polyclonal Antibody

Catalog # AP72081

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q03431</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	66361

## Additional Information

---

<b>Gene ID</b>	5745
<b>Other Names</b>	PTH1R; PTHR; PTHR1; Parathyroid hormone/parathyroid hormone-related peptide receptor; PTH/PTHrP type I receptor; PTH/PTHr receptor; Parathyroid hormone 1 receptor; PTH1 receptor
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	PTH1R {ECO:0000303   PubMed:10913300, ECO:0000312   HGNC:HGNC:9608}
<b>Function</b>	G-protein-coupled receptor for parathyroid hormone (PTH) and for parathyroid hormone-related peptide (PTHLH) (PubMed: <a href="#">10913300</a> , PubMed: <a href="#">18375760</a> , PubMed: <a href="#">19674967</a> , PubMed: <a href="#">27160269</a> , PubMed: <a href="#">30975883</a> , PubMed: <a href="#">35932760</a> , PubMed: <a href="#">8397094</a> ). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (cAMP) (PubMed: <a href="#">30975883</a> , PubMed: <a href="#">35932760</a> ). PTH1R is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed: <a href="#">20172855</a> , PubMed: <a href="#">30975883</a> , PubMed: <a href="#">35932760</a> ). PTHLH dissociates from PTH1R more rapidly than PTH; as consequence, the cAMP response induced by PTHLH decays faster than the response induced by PTH (PubMed: <a href="#">35932760</a> ).
<b>Cellular Location</b>	Cell membrane; Multi-pass membrane protein


## Tissue Location

Expressed in most tissues. Most abundant in kidney, bone and liver.

## Background

Receptor for parathyroid hormone and for parathyroid hormone-related peptide. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and also a phosphatidylinositol-calcium second messenger system.

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.