

ADRB2 Antibody (S364)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7263E

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P07550
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB16260
Calculated MW	46459
Antigen Region	345-373

Additional Information

Gene ID	154
Other Names	Beta-2 adrenergic receptor, Beta-2 adrenoreceptor, Beta-2 adrenoceptor, ADRB2, ADRB2R, B2AR
Target/Specificity	This ADRB2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 345-373 amino acids from human ADRB2.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ADRB2 Antibody (S364) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ADRB2 (HGNC:286)
Synonyms	ADRB2R, B2AR
Function	G protein-coupled receptor for catecholamines that couples to both G(s)

and G(i) proteins, activating bifurcated signaling pathways (PubMed:[2831218](#), PubMed:[7915137](#)). ADRB2 binds epinephrine (Epi) with an approximately 30-fold greater affinity than norepinephrine (NE) (PubMed:[2831218](#), PubMed:[33093660](#), PubMed:[7915137](#)). In the heart, Epi- and NE-activated ADRB2 induces rapid and slow cardiomyocyte contraction rate, respectively (By similarity). Both NE and Epi promote coupling to G(s)/PKA pathway to regulate myocyte contraction rate (By similarity). Epi also promotes ADRB2 coupling to G(i) proteins to exert cardioprotective effects especially in the conditions of hypoxia and oxidative stress through the G(i)/PI3K/Akt signaling pathway (By similarity). ADRB2-G(s) signaling delivers proapoptotic signals in cardiomyocytes although G(i)-mediated survival effect appears to predominate (By similarity). ADRB2 also transduces signals independently of PKA to regulate cellular pH by modulating Na(+)/H(+) exchanger SLC9A3 function (PubMed:[9560162](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein. Golgi apparatus. Note=Colocalizes with VHL at the cell membrane (PubMed:19584355). Activated receptors are internalized into endosomes prior to their degradation in lysosomes (PubMed:20559325). Activated receptors are also detected within the Golgi apparatus (PubMed:27481942).


Background

ADRB2, beta-2-adrenergic receptor which is a member of the G protein-coupled receptor superfamily. This receptor is directly associated with one of its ultimate effectors, the class C L-type calcium channel Ca(V)1.2. This receptor-channel complex also contains a G protein, an adenylyl cyclase, cAMP-dependent kinase, and the counterbalancing phosphatase, PP2A. The assembly of the signaling complex provides a mechanism that ensures specific and rapid signaling by this G protein-coupled receptor. This protein is intronless.

References


Kobilka B.K., Dixon R.A.F. Proc. Natl. Acad. Sci. U.S.A. 84:46-50(1987)
Emorine L.J., Marullo S. Proc. Natl. Acad. Sci. U.S.A. 84:6995-6999(1987)
Kobilka B.K., Frielle T.J. Biol. Chem. 262:7321-7327(1987)

Images




Immunohistochemical analysis of AP7263E on paraffin-embedded Human hepato carcinoma tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

Overlay histogram showing A431 cells stained with AP7263E(green line). The cells were fixed with 2% paraformaldehyde and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody



(1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1 μ g/1 \times 10 6 cells) used under the same conditions. Acquisition of >10,000 events was performed.



All lanes : Anti-ADRB2 Antibody (S364) at 1:2000 dilution
 Lane 1: A431 whole cell lysate Lane 2: PC-3 whole cell lysate Lane 3: HUVEC whole cell lysate Lane 4: HeLa whole cell lysate Lane 5: Human liver lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 46 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.