

# MMP-3 Polyclonal Antibody

Catalog # AP72648

## Product Information

---

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P08254</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53977

## Additional Information

---

Gene ID	4314
Other Names	MMP3; STMY1; Stromelysin-1; SL-1; Matrix metalloproteinase-3; MMP-3; Transin-1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	MMP3
Synonyms	STMY1
Function	Metalloproteinase with a rather broad substrate specificity that can degrade fibronectin, laminin, gelatins of type I, III, IV, and V; collagens III, IV, X, and IX, and cartilage proteoglycans. Activates different molecules including growth factors, plasminogen or other matrix metalloproteinases such as MMP9 (PubMed: <a href="#">11029580</a> , PubMed: <a href="#">1371271</a> ). Once released into the extracellular matrix (ECM), the inactive pro-enzyme is activated by the plasmin cascade signaling pathway (PubMed: <a href="#">2383557</a> ). Also acts intracellularly (PubMed: <a href="#">22265821</a> ). For example, in dopaminergic neurons, gets activated by the serine protease HTRA2 upon stress and plays a pivotal role in DA neuronal degeneration by mediating microglial activation and alpha- synuclein/SNCA cleavage (PubMed: <a href="#">21330369</a> ). In addition, plays a role in immune response and possesses antiviral activity against various viruses such as vesicular stomatitis virus, influenza A virus (H1N1) and human herpes virus 1 (PubMed: <a href="#">35940311</a> ). Mechanistically, translocates from the cytoplasm into

the cell nucleus upon virus infection to influence NF-kappa-B activities (PubMed:[35940311](#)).


## Cellular Location

Secreted, extracellular space, extracellular matrix. Nucleus. Cytoplasm


## Background

Can degrade fibronectin, laminin, gelatins of type I, III, IV, and V; collagens III, IV, X, and IX, and cartilage proteoglycans. Activates procollagenase.

## Images



Western Blot analysis of various cells using MMP-3 Polyclonal Antibody diluted at 1 : 2000



Western Blot analysis of 293 cells using MMP-3 Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.