

# TRAP220 Polyclonal Antibody

Catalog # AP72907

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q15648</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	168478

## Additional Information

---

<b>Gene ID</b>	5469
<b>Other Names</b>	MED1; ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; Mediator of RNA polymerase II transcription subunit 1; Activator-recruited cofactor 205 kDa component; ARC205; Mediator complex subunit 1; Peroxiso
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	MED1
<b>Function</b>	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene- specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (PubMed: <a href="#">10406464</a> , PubMed: <a href="#">11867769</a> , PubMed: <a href="#">12037571</a> , PubMed: <a href="#">12218053</a> , PubMed: <a href="#">12556447</a> , PubMed: <a href="#">14636573</a> , PubMed: <a href="#">15340084</a> , PubMed: <a href="#">15471764</a> , PubMed: <a href="#">15989967</a> , PubMed: <a href="#">16574658</a> , PubMed: <a href="#">9653119</a> ). Acts as a coactivator for GATA1-mediated transcriptional activation during erythroid differentiation of K562 erythroleukemia cells (PubMed: <a href="#">24245781</a> ).

**Cellular Location** Nucleus. Note=A subset of the protein may enter the nucleolus subsequent to phosphorylation by MAPK1 or MAPK3

**Tissue Location** Ubiquitously expressed.


## Background

---

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (PubMed:[10406464](#), PubMed:[11867769](#), PubMed:[12037571](#), PubMed:[12218053](#), PubMed:[12556447](#), PubMed:[14636573](#), PubMed:[15340084](#), PubMed:[15471764](#), PubMed:[15989967](#), PubMed:[16574658](#), PubMed:[9653119](#)). Acts as a coactivator for GATA1-mediated transcriptional activation during erythroid differentiation of K562 erythroleukemia cells (PubMed:[24245781](#)).

## Images

---



Western Blot analysis of various cells using TRAP220 Polyclonal Antibody diluted at 1 : 500. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.