

USP30 Polyclonal Antibody

Catalog # AP73017

Product Information

Application	WB, E
Primary Accession	Q70CQ3
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58503

Additional Information

Gene ID	84749
Other Names	USP30; Ubiquitin carboxyl-terminal hydrolase 30; Deubiquitinating enzyme 30; Ubiquitin thioesterase 30; Ubiquitin-specific-processing protease 30; Ub-specific protease 30
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	USP30 (HGNC:20065)
Function	Deubiquitinating enzyme tethered to the mitochondrial outer membrane that acts as a key inhibitor of mitophagy by counteracting the action of parkin (PRKN): hydrolyzes ubiquitin attached by parkin on target proteins, such as RHOT1/MIRO1 and TOMM20, thereby blocking parkin's ability to drive mitophagy (PubMed: 18287522 , PubMed: 24896179 , PubMed: 25527291 , PubMed: 25621951). Preferentially cleaves 'Lys-6'- and 'Lys-11'-linked polyubiquitin chains, 2 types of linkage that participate in mitophagic signaling (PubMed: 25621951). Does not cleave efficiently polyubiquitin phosphorylated at 'Ser-65' (PubMed: 25527291). Acts as a negative regulator of mitochondrial fusion by mediating deubiquitination of MFN1 and MFN2 (By similarity).
Cellular Location	Mitochondrion outer membrane
Tissue Location	Expressed in skeletal muscle, pancreas, liver and kidney.

Background

Deubiquitinating enzyme tethered to the mitochondrial outer membrane that acts as a key inhibitor of mitophagy by counteracting the action of parkin (PRKN): hydrolyzes ubiquitin attached by parkin on target proteins, such as RHOT1/MIRO1 and TOMM20, thereby blocking parkin's ability to drive mitophagy (PubMed:[18287522](#), PubMed:[24896179](#), PubMed:[25527291](#), PubMed:[25621951](#)). Preferentially cleaves 'Lys-6'- and 'Lys-11'- linked polyubiquitin chains, 2 types of linkage that participate to mitophagic signaling (PubMed:[25621951](#)). Does not cleave efficiently polyubiquitin phosphorylated at 'Ser-65' (PubMed:[25527291](#)). Acts as negative regulator of mitochondrial fusion by mediating deubiquitination of MFN1 and MFN2 (By similarity).

Images



Western Blot analysis of various cells using USP30 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.