

β-1,4-Gal-T5 Polyclonal Antibody

Catalog # AP73193

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, E |
| Primary Accession | O43286 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 45119 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 9334 |
| Other Names | B4GALT5; Beta-1; 4-galactosyltransferase 5; Beta-1, 4-GalTase 5; Beta4Gal-T5; b4Gal-T5; Beta-1, 4-GALT II; UDP-Gal:beta-GlcNAc beta-1, 4-galactosyltransferase 5; UDP-galactose:beta-N-acetylglucosamine beta-1, 4-galactosyltransferase 5 |
| Dilution | WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A |
| Format | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide. |
| Storage Conditions | -20°C |


Protein Information

| | |
|--------------------------|---|
| Name | B4GALT5 (HGNC:928) |
| Function | Catalyzes the synthesis of lactosylceramide (LacCer) via the transfer of galactose from UDP-galactose to glucosylceramide (GlcCer) (PubMed: 24498430). LacCer is the starting point in the biosynthesis of all gangliosides (membrane-bound glycosphingolipids) which play pivotal roles in the CNS including neuronal maturation and axonal and myelin formation (By similarity). Plays a role in the glycosylation of BMPR1A and regulation of its protein stability (By similarity). Essential for extraembryonic development during early embryogenesis (By similarity). |
| Cellular Location | Golgi apparatus, Golgi stack membrane {ECO:0000250 UniProtKB:P15291}; Single-pass type II membrane protein Golgi apparatus {ECO:0000250 UniProtKB:A0A1S6M251}. Note=Trans cisternae of Golgi stack. {ECO:0000250 UniProtKB:P15291} |
| Tissue Location | Ubiquitously expressed. |


Background

Catalyzes the synthesis of lactosylceramide (LacCer) via the transfer of galactose from UDP-galactose to glucosylceramide (GlcCer) (PubMed:[24498430](#)). LacCer is the starting point in the biosynthesis of all gangliosides (membrane-bound glycosphingolipids) which play pivotal roles in the CNS including neuronal maturation and axonal and myelin formation (By similarity). Plays a role in the glycosylation of BMPR1A and regulation of its protein stability (By similarity). Essential for extraembryonic development during early embryogenesis (By similarity).

Images



Western Blot analysis of various cells using β -1,4-Gal-T5 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000



Western Blot analysis of 293 cells using β -1,4-Gal-T5 Polyclonal Antibody diluted at 1 : 1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.