

# FGG Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP7325b

## Product Information

---

<b>Application</b>	IHC-P, FC, WB, E
<b>Primary Accession</b>	<a href="#">P02679</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	51512
<b>Antigen Region</b>	417-445

## Additional Information

---

<b>Gene ID</b>	2266
<b>Other Names</b>	Fibrinogen gamma chain, FGG
<b>Target/Specificity</b>	This FGG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 417-445 amino acids from the C-terminal region of human FGG.
<b>Dilution</b>	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	FGG Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FGG
<b>Function</b>	Together with fibrinogen alpha (FGA) and fibrinogen beta (FGB), polymerizes to form an insoluble fibrin matrix. Has a major function in hemostasis as one of the primary components of blood clots. In addition, functions during the early stages of wound repair to stabilize the lesion and guide cell migration during re-epithelialization. Was originally thought to be essential for platelet

aggregation, based on in vitro studies using anticoagulated blood. However, subsequent studies have shown that it is not absolutely required for thrombus formation in vivo. Enhances expression of SELP in activated platelets via an ITGB3-dependent pathway. Maternal fibrinogen is essential for successful pregnancy. Fibrin deposition is also associated with infection, where it protects against IFNG-mediated hemorrhage. May also facilitate the antibacterial immune response via both innate and T-cell mediated pathways.

<b>Cellular Location</b>	Secreted
<b>Tissue Location</b>	Detected in blood plasma (at protein level).


## Background

FGG is the gamma component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this protein lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia and thrombophilia.


## References

- Sie, M.P., Isaacs, A. J. *Hypertens.* 27 (7), 1392-1398 (2009)  
Nowak-Gottl, U., Weiler, H. *Blood* (2009) In press  
de Willige, S.U., Pyle, M.E. *Thromb. Haemost.* 101 (6), 1078-1084 (2009)  
Yoshida, N., Imaoka, S. *Thromb. Haemost.* 68 (5), 534-538 (1992)


## Images



FGG Antibody (C-term) (Cat. #AP7325b) western blot analysis in human blood plasma cell line lysates (35ug/lane). This demonstrates the FGG antibody detected the FGG protein (arrow).



FGG Antibody (C-term) (RB18714) IHC analysis in formalin fixed and paraffin embedded human hepatocarcinoma tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FGG Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



FGG Antibody (C-term) (Cat. #AP7325b) flow cytometric analysis of T47D cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.