

# CA IV Polyclonal Antibody

Catalog # AP73311

## Product Information

---

Application	WB
Primary Accession	<a href="#">P22748</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35032

## Additional Information

---

Gene ID	762
Other Names	CA4; Carbonic anhydrase 4; Carbonate dehydratase IV; Carbonic anhydrase IV; CA-IV
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	CA4 ( <a href="#">HGNC:1375</a> )
Function	Catalyzes the reversible hydration of carbon dioxide into bicarbonate and protons and thus is essential to maintaining intracellular and extracellular pH (PubMed: <a href="#">15563508</a> , PubMed: <a href="#">16686544</a> , PubMed: <a href="#">16807956</a> , PubMed: <a href="#">17127057</a> , PubMed: <a href="#">17314045</a> , PubMed: <a href="#">17652713</a> , PubMed: <a href="#">17705204</a> , PubMed: <a href="#">18618712</a> , PubMed: <a href="#">19186056</a> , PubMed: <a href="#">19206230</a> , PubMed: <a href="#">7625839</a> ). May stimulate the sodium/bicarbonate transporter activity of SLC4A4 that acts in pH homeostasis (PubMed: <a href="#">15563508</a> ). It is essential for acid overload removal from the retina and retina epithelium, and acid release in the choriocapillaris in the choroid (PubMed: <a href="#">15563508</a> ).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor
Tissue Location	Expressed in the endothelium of the choriocapillaris in eyes (at protein level). Not expressed in the retinal epithelium at detectable levels.


## Background

---

Reversible hydration of carbon dioxide. May stimulate the sodium/bicarbonate transporter activity of SLC4A4 that acts in pH homeostasis. It is essential for acid overload removal from the retina and retina epithelium, and acid release in the choriocapillaris in the choroid.

## Images

---



Western Blot analysis of rat muscle cells using CA IV Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.