

# Fascin 1 Polyclonal Antibody

Catalog # AP73395

# **Product Information**

**Application** WB, IHC-P **Primary Accession** 016658

**Reactivity** Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW54530

### **Additional Information**

Gene ID 6624

Other Names FSCN1; FAN1; HSN; SNL; Fascin; 55 kDa actin-bundling protein; Singed-like

protein; p55

**Dilution** WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet

tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p:

1:100-300 ELISA: 1/20000. Not yet tested in other applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name FSCN1

**Synonyms** FAN1, HSN, SNL

**Function** Actin-binding protein that contains 2 major actin binding sites

(PubMed:<u>21685497</u>, PubMed:<u>23184945</u>). Organizes filamentous actin into parallel bundles (PubMed:<u>20393565</u>, PubMed:<u>21685497</u>, PubMed:<u>23184945</u>). Plays a role in the organization of actin filament bundles and the formation of microspikes, membrane ruffles, and stress fibers (PubMed:<u>22155786</u>). Important for the formation of a diverse set of cell protrusions, such as

filopodia, and for cell motility and migration (PubMed: 20393565,

PubMed: <u>21685497</u>, PubMed: <u>23184945</u>). Mediates reorganization of the actin

cytoskeleton and axon growth cone collapse in response to NGF

(PubMed:22155786).

**Cellular Location** Cytoplasm, cytosol. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton.

Cytoplasm, cytoskeleton, stress fiber. Cell projection, filopodium. Cell projection, invadopodium. Cell projection, microvillus. Cell junction.

Note=Colocalized with RUFY3 and F-actin at filipodia of the axonal growth cone. Colocalized with DBN1 and F- actin at the transitional domain of the axonal growth cone (By similarity). {ECO:0000250|UniProtKB:Q61553, ECO:0000269|PubMed:21706053}


**Tissue Location** 

Ubiquitous.

# **Background**

Organizes filamentous actin into bundles with a minimum of 4.1:1 actin/fascin ratio. Plays a role in the organization of actin filament bundles and the formation of microspikes, membrane ruffles, and stress fibers. Important for the formation of a diverse set of cell protrusions, such as filopodia, and for cell motility and migration.


## **Images**



Western Blot analysis of A549, HuvEc, HT29, K562, L929 cells using Fascin 1 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.