

CD39 Polyclonal Antibody

Catalog # AP73769

Product Information

Application	WB, E
Primary Accession	P49961
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57965

Additional Information

Gene ID	953
Other Names	ENTPD1; CD39; Ectonucleoside triphosphate diphosphohydrolase 1; NTPDase 1; Ecto-ATP diphosphohydrolase 1; Ecto-ATPDase 1; Ecto-ATPase 1; Ecto-apyrase; Lymphoid cell activation antigen; CD39
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	ENTPD1 (HGNC:3363)
Function	Catalyzes the hydrolysis of nucleoside triphosphates (NTPs) and diphosphates (NDPs) (Probable) (PubMed: 8529670 , PubMed: 8626624 , PubMed: 8955160 , PubMed: 8996251). The enzyme sequentially removes phosphate groups in two successive steps, converting NTPs to nucleoside monophosphates (NMPs) via NDP intermediates (Probable) (PubMed: 8529670 , PubMed: 8626624 , PubMed: 8955160 , PubMed: 8996251). This activity contributes to the regulation of extracellular levels of nucleotides (Probable) (PubMed: 8529670 , PubMed: 8626624 , PubMed: 8955160 , PubMed: 8996251). By hydrolyzing proinflammatory ATP and platelet-activating ADP to AMP, it blocks platelet aggregation and supports blood flow (PubMed: 8955160 , PubMed: 8996251).
Cellular Location	Membrane; Multi-pass membrane protein. Membrane, caveola
Tissue Location	Expressed primarily on activated lymphoid cells (PubMed:7930580). Also expressed in endothelial tissues (PubMed:8955160). Highly expressed in

placenta, lung, skeletal muscle, kidney (PubMed:8955160).

Background

In the nervous system, could hydrolyze ATP and other nucleotides to regulate purinergic neurotransmission. Could also be implicated in the prevention of platelet aggregation by hydrolyzing platelet-activating ADP to AMP. Hydrolyzes ATP and ADP equally well.

Images



Western Blot analysis of HeLa cells using CD39 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.