

CD315 Polyclonal Antibody

Catalog # AP73820

Product Information

Application WB, IHC-P, IF, ICC, E

Q9P2B2 Primary Accession

Human, Mouse, Rat Reactivity

Host Rabbit Clonality Polyclonal Calculated MW 98556

Additional Information

Gene ID 5738

Other Names PTGFRN; CD9P1; EWIF; FPRP; KIAA1436; Prostaglandin F2 receptor negative

> regulator; CD9 partner 1; CD9P-1; Glu-Trp-Ile EWI motif-containing protein F; EWI-F; Prostaglandin F2-alpha receptor regulatory protein; Prostaglandin

F2-alpha receptor-associated protein; CD315

Dilution WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not

yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. IHC-p:

1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

IF~~1:50~200 ICC~~N/A E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name **PTGFRN**

Synonyms CD9P1, EWIF, FPRP, KIAA1436

Function Inhibits the binding of prostaglandin F2-alpha (PGF2-alpha) to its specific FP

> receptor, by decreasing the receptor number rather than the affinity constant. Functional coupling with the prostaglandin F2-alpha receptor seems to occur (By similarity). In myoblasts, associates with tetraspanins CD9 and CD81 to

prevent myotube fusion during muscle regeneration (By similarity).

Cellular Location Endoplasmic reticulum membrane; Single-pass type I membrane protein.


Golgi apparatus, trans-Golgi network membrane; Single-pass type I

membrane protein


Background

Inhibits the binding of prostaglandin F2-alpha (PGF2- alpha) to its specific FP receptor, by decreasing the receptor number rather than the affinity constant. Functional coupling with the prostaglandin F2-alpha receptor seems to occur (By similarity).

Images



Western Blot analysis of 293 cells using CD315 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.