

HLA-DMβ Polyclonal Antibody

Catalog # AP74208

Product Information

Application IHC-P
Primary Accession P28068
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 28931

Additional Information

Gene ID 3109

Other Names HLA class II histocompatibility antigen, DM beta chain (MHC class II antigen

DMB) (Really interesting new gene 7 protein)

Dilution IHC-P~~IHC-p 1:50-200, ELISA 1:10000-20000

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name HLA-DMB

Synonyms DMB, RING7

Function Plays a critical role in catalyzing the release of class II- associated invariant

chain peptide (CLIP) from newly synthesized MHC class II molecules and freeing the peptide binding site for acquisition of antigenic peptides. In B-cells, the interaction between HLA-DM and MHC class II molecules is

regulated by HLA-DO.

Cellular Location Late endosome membrane; Single-pass type I membrane protein. Lysosome


membrane; Single-pass type I membrane protein. Note=Localizes to late

endocytic compartment. Associates with lysosome membranes


Background

Plays a critical role in catalyzing the release of class II-associated invariant chain peptide (CLIP) from newly synthesized MHC class II molecules and freeing the peptide binding site for acquisition of antigenic peptides. In B-cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO.


Images



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.