

TNFAIP8L3 Polyclonal Antibody

Catalog # AP74255

Product Information

Application	WB
Primary Accession	Q5GJ75
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32659

Additional Information

Gene ID	388121
Other Names	Tumor necrosis factor alpha-induced protein 8-like protein 3 (TNF alpha-induced protein 8-like protein 3) (TNFAIP8-like protein 3)
Dilution	WB~~WB 1:500-2000, ELISA 1:10000-20000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	TNFAIP8L3
Synonyms	TIPE3
Function	Acts as a lipid transfer protein. Preferentially captures and shuttles two lipid second messengers, i.e., phosphatidylinositol 4,5- biphosphate and phosphatidylinositol 3,4,5-trisphosphate and increases their levels in the plasma membrane. Additionally, may also function as a lipid-presenting protein to enhance the activity of the PI3K-AKT and MEK-ERK pathways. May act as a regulator of tumorigenesis through its activation of phospholipid signaling.
Cellular Location	Cytoplasm. Cell membrane {ECO:0000250 UniProtKB:Q3TBL6}. Note=On PDGF activation, translocates from cytoplasm to plasma membrane {ECO:0000250 UniProtKB:Q3TBL6}
Tissue Location	Widely expressed (at protein level) (PubMed:25479791). Highly expressed in most carcinoma cell lines (PubMed:25242044, PubMed:25479791).

Background

Acts as a lipid transfer protein. Preferentially captures and shuttles two lipid second messengers, i.e., phosphatidylinositol 4,5- bisphosphate and phosphatidylinositol 3,4,5-trisphosphate and increases their levels in the plasma membrane. Additionally, may also function as a lipid-presenting protein to enhance the activity of the PI3K-AKT and MEK-ERK pathways. May act as a regulator of tumorigenesis through its activation of phospholipid signaling.

Images



Western Blot analysis of 1,293T 2, mouse-brain cells using primary antibody diluted at 1:1000(4°C overnight).
Secondary antibody : Goat Anti-rabbit IgG IRDye 800(diluted at 1:5000, 25°C, 1 hour)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.