

Rab5C Rabbit mAb

Catalog # AP74883

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	P51148
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	23483

Additional Information

Gene ID	5878
Other Names	RAB5C
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IHC-F~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	RAB5C (HGNC:9785)
Synonyms	RABL
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed: 16086013 , PubMed: 17562788). Involved in early endocytic trafficking (PubMed: 16086013 , PubMed: 17562788). Required for EEA1 recruitment to early endosomes (PubMed: 16086013 , PubMed: 17562788). Required for EGF and transferrin endocytosis and trafficking through early endosomes (PubMed: 16086013 , PubMed: 17562788).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P20339}; Lipid-anchor {ECO:0000250 UniProtKB:P20339}; Cytoplasmic side

{ECO:0000250|UniProtKB:P20339}. Early endosome membrane
{ECO:0000250|UniProtKB:P20339}; Lipid-anchor
{ECO:0000250|UniProtKB:P20339}. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV


Background

Protein transport.


Images




Immunocytochemistry analysis of Rab5C (green) in A549 using Rab5C antibody, and DAPI (blue).



Western blot analysis of RAB5C in K562, C6, 3T3, HeLa lysates using RAB5C antibody.



Immunohistochemistry analysis of paraffin-embedded Human Cholangiocarcinoma using RAB5C antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.