

Lactate Dehydrogenase A Rabbit mAb

Catalog # AP74949

Product Information

Application	WB, IHC-P, IHC-F, FC, IP
Primary Accession	P00338
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	36689

Additional Information

Gene ID	3939
Other Names	LDHA
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IHC-F~~N/A FC~~1:50-1:100 IP~~1:20-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	LDHA (HGNC:6535)
Function	Interconverts simultaneously and stereospecifically pyruvate and lactate with concomitant interconversion of NADH and NAD(+).
Cellular Location	Cytoplasm.
Tissue Location	Predominantly expressed in anaerobic tissues such as skeletal muscle and liver.





Background


Lactate dehydrogenase (LDH) catalyzes the interconversion of pyruvate and NADH to lactate and NAD⁺. When the oxygen supply is too low for mitochondrial ATP production, this reaction recycles NADH generated in glycolysis to NAD⁺, which reenters glycolysis. The major form of LDH found in muscle cells is the A (LDHA)

isozyme.

Images

Western blot analysis of LDHA in Jurkat, rat Brain, C6, HeLa lysates using Lactate Dehydrogenase A antibody.





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.