

# GIPR Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7495A

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P48546</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	53157
<b>Antigen Region</b>	7-38

## Additional Information

---

<b>Gene ID</b>	2696
<b>Other Names</b>	Gastric inhibitory polypeptide receptor, GIP-R, Glucose-dependent insulinotropic polypeptide receptor, GIPR
<b>Target/Specificity</b>	This GIPR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-38 amino acids from the N-terminal region of human GIPR.
<b>Dilution</b>	WB~~1:500 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	GIPR Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GIPR
<b>Function</b>	This is a receptor for GIP. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase.
<b>Cellular Location</b>	Cell membrane; Multi-pass membrane protein


## Background

GIPR also called glucose-dependent insulinotropic polypeptide, is a 42-amino acid polypeptide synthesized by K cells of the duodenum and small intestine. This protein was originally identified as an activity in gut extracts that inhibited gastric acid secretion and gastrin release, but subsequently was demonstrated to stimulate insulin release potently in the presence of elevated glucose. The insulinotropic effect on pancreatic islet beta-cells was then recognized to be the principal physiologic action of GIP. Together with glucagon-like peptide-1, GIP is largely responsible for the secretion of insulin after eating. The protein is involved in several other facets of the anabolic response.


## References

Herbach,N. Am. J. Physiol. Renal Physiol. 296 (4), F819-F829 (2009) Rudovich,N., Kaiser,S. Regul. Pept. 142 (3), 138-145 (2007) Nitz,I., Fisher,E. Mol Nutr Food Res 51 (8), 1046-1052 (2007)


## Images




All lanes : Anti-GIPR Antibody (N-term) at 1:1000 dilution  
Lane 1: HL-60 whole cell lysate Lane 2: R. brain whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 53kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of GIPR antibody (N-term) (Cat.#AP7495a) in HL60 cell line lysates (35ug/lane). GIPR (arrow) was detected using the purified Pab.



GIPR Antibody (N-term) (Cat. #AP7495A) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GIPR Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



GIPR Antibody (Center) (Cat. #AP7495a) flow cytometric analysis of MDA-MB231 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## Citations

---

- [Transgenic rescue of adipocyte glucose-dependent insulinotropic polypeptide receptor expression restores high fat diet-induced body weight gain.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.