

# ApoER2 Rabbit mAb

Catalog # AP75092

## Product Information

---

Application	WB, IP
Primary Accession	<a href="#">Q14114</a>
Reactivity	Human, Rat, Hamster
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	105634

## Additional Information

---

Gene ID	7804
Other Names	LRP8
Dilution	WB~~1/500-1/1000 IP~~1/20
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

Name	LRP8
Synonyms	APOER2
Function	Cell surface receptor for Reelin (RELN) and apolipoprotein E (apoE)-containing ligands (PubMed: <a href="#">12899622</a> , PubMed: <a href="#">12950167</a> , PubMed: <a href="#">20223215</a> , PubMed: <a href="#">30873003</a> ). LRP8 participates in transmitting the extracellular Reelin signal to intracellular signaling processes, by binding to DAB1 on its cytoplasmic tail (By similarity). Reelin acts via both the VLDL receptor (VLDLR) and LRP8 to regulate DAB1 tyrosine phosphorylation and microtubule function in neurons (By similarity). LRP8 has higher affinity for Reelin than VLDLR (By similarity). LRP8 is thus a key component of the Reelin pathway which governs neuronal layering of the forebrain during embryonic brain development (By similarity). Binds the endoplasmic reticulum resident receptor- associated protein (RAP) (By similarity). Binds dimers of beta 2-glycoprotein I and may be involved in the suppression of platelet aggregation in the vasculature (PubMed: <a href="#">12807892</a> ). Highly expressed in the initial segment of the epididymis, where it affects the functional expression of clusterin and phospholipid hydroperoxide glutathione peroxidase (PHGPx), two proteins required for sperm maturation (By similarity). May also function

as an endocytic receptor (By similarity). Not required for endocytic uptake of SEPP1 in the kidney which is mediated by LRP2 (By similarity). Together with its ligand, apolipoprotein E (apoE), may indirectly play a role in the suppression of the innate immune response by controlling the survival of myeloid- derived suppressor cells (By similarity).

### Cellular Location


Cell membrane; Single-pass type I membrane protein. Secreted {ECO:0000250|UniProtKB:Q924X6}. Note=Isoforms that contain the exon coding for a furin-type cleavage site are proteolytically processed, leading to a secreted receptor fragment {ECO:0000250|UniProtKB:Q924X6}

### Tissue Location

Expressed mainly in brain and placenta. Also expressed in platelets and megakaryocytic cells. Not expressed in the liver.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.