

# Aquaporin 5 Rabbit mAb

Catalog # AP75098

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, ICC
<b>Primary Accession</b>	<a href="#">P55064</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Calculated MW</b>	28292

## Additional Information

---

<b>Gene ID</b>	362
<b>Other Names</b>	AQP5
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A ICC~~N/A
<b>Format</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


## Protein Information

---

<b>Name</b>	AQP5 ( <a href="#">HGNC:638</a> )
<b>Function</b>	Aquaporins form homotetrameric transmembrane channels, with each monomer independently mediating water transport across the plasma membrane along its osmotic gradient (PubMed: <a href="#">18768791</a> , PubMed: <a href="#">8621489</a> ). Plays an important role in fluid secretion in salivary glands (By similarity). Required for TRPV4 activation by hypotonicity. Together with TRPV4, controls regulatory volume decrease in salivary epithelial cells (PubMed: <a href="#">16571723</a> ). Seems to play a redundant role in water transport in the eye, lung and in sweat glands (By similarity).
<b>Cellular Location</b>	Apical cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein Note=Hypotonicity increases location at the cell membrane Phosphorylation decreases location at the cell membrane
<b>Tissue Location</b>	Detected in skin eccrine sweat glands, at the apical cell membrane and at intercellular canaliculi (at protein level).

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.